arXiv:1610.10043v1 [math.KT] 31 Oct 2016

Infinity categories with duality and hermitian multiplicative infinite loop space machines

Hadrian Heine, Alejo Lopez-Avila, Markus Spitzweck

November 1, 2016

Abstract

We show that any symmetric monoidal ∞-category with duality gives rise to a direct sum hermitian K-theory spectrum. This assignment is lax symmetric monoidal, thereby producing E_∞-ring spectra from bimonoidal ∞-categories with duality.

To produce examples of such structures we give two hermitian recognition principles. Combining these we show that any rigid symmetric monoidal ∞-category which is preadditive gives rise to a bimonoidal ∞-category with duality.

We twist the dualities in various ways and apply our definitions to finitely generated projective modules over E_∞-ring spectra.

Contents

1 Introduction **2**

2 ∞-categorical background **3**

3 Infinity categories with duality **5**

4 Symmetric monoidal structures **5**

5 Group completions and spectra **8**

6 The first hermitian recognition principle **9**

7 The second hermitian recognition principle **12**

8 The extended second hermitian recognition principle **17**

9 Twisting the duality **24**

10 C_2-equivariant K-theory **31**

11 Hermitian objects **32**

12 Applications **34**

1 Introduction

K-theory has by now a longstanding history. Vector bundles on topological spaces or schemes give rise to the K_0-functor and also higher K-theory ([12]). Algebraic K-theory can be defined for any stable ∞-category ([3]) or Waldhausen (∞-) category ([19], [1]).

Direct sum K-theory of a symmetric monoidal ∞-category C can be defined to be the spectrum $K(C)$ associated to the group completion of the E_∞-space $C^{\sim}$, the underlying ∞-groupoid of C together with its symmetric monoidal structure ([4]). It was shown in loc. cit. that the functor K is in fact lax symmetric monoidal, using a symmetric monoidal structure on the ∞-category of small symmetric monoidal ∞-categories resembling the one on E_∞-spaces (defined in [11] for Γ-spaces and studied in [16]).

Instead of vector bundles one can consider vector bundles equiped with non-degenerate symmetric or anti-symmetric bilinear forms giving rise to the Grothendieck-Witt group and hermitian K-theory. Motivated by direct sum K-theory we consider in this article a version of direct sum hermitian K-theory. Given a symmetric monoidal category C with duality, e.g. finitely generated projective R-modules $\mathcal{P}(R)$ over a commutative ring R with the direct sum as symmetric monoidal structure and the usual duality, one can consider the symmetric monoidal groupoid of the symmetric monoidal category C_h of hermitian objects of C and the spectrum associated to its group completion. This version of hermitian K-theory coincides with the common definitions in the case of $\mathcal{P}(R)$ (see [14] and [13, Theorem A.1] if 2 is invertible in R and [5, Theorem A] in general, compare also to [6, 1.6. Theoreme]).

Direct sum K-theory is also meaningful in the case of modules over E_∞-ring spectra, e.g. the direct sum K-theory of finitely generated projective R-modules for R a connective E_∞-ring spectrum is the connective algebraic K-theory of R ([8, Theorem 5]).

We expect a similar behavior in the case of hermitian K-theory. Hermitian K-theory via a Waldhausen type construction is not developed for E_∞-ring spectra so far. We partially remedy this defect in this article by exhibiting perfect or finitely generated projective modules over an E_∞-ring spectrum as an ∞-category with duality together with the relevant symmetric monoidal structures, to which direct sum hermitian K-theory can be applied, thereby producing an E_∞-ring spectrum.

Our definition of an ∞-category with duality is fairly straight forward, it is a homotopy fixed point of the action of the group on two elements C_2 on the ∞-category of small ∞-categories Cat_∞, where the non-trivial element of C_2 sends a $C \in \mathsf{Cat}_\infty$ to C^{op}.

We show that this definition is equivalent to giving on a $C \in \mathsf{Cat}_\infty$ a symmetric perfect pairing $C \times C \to \mathsf{Spc}$, where Spc denotes the ∞-category of spaces, see Corollary 7.3.

We equip the ∞-category of small symmetric monoidal ∞-categories with duality, which will be denoted by $\mathsf{SymMonCat}_\infty^{hC_2}$, with a symmetric monoidal structure (section 4) and show that the associated hermitian K-theory functor is lax symmetric monoidal with respect to this symmetric monoidal structure (Proposition 5.1).

Our first hermitian recognition principle exhibits every symmetric monoidal ∞-category C with duality which is preadditive as a commutative monoid object

in $\mathsf{SymMonCat}_\infty^{hC_2}$, where morally the given symmetric monoidal structure on C will give rise to the second symmetric monoidal structure and the coproduct on C to the first symmetric monoidal structure which a commutative monoid in $\mathsf{SymMonCat}_\infty^{hC_2}$ carries.

The second hermitian recognition principle provides for any rigid symmetric monoidal ∞-category C an object of $\mathsf{SymMonCat}_\infty^{hC_2}$. Morally the duality is given by sending an object $X \in C$ to the dual object $X^\vee$. The non-trivial part to show is that this really exhibits a duality on C with all higher coherences and that the duality is in fact compatible with the symmetric monoidal structure. This is the content of section 7.

Section 8 extends the results of section 7 to the case where the duality is twisted by a tensor invertible object in the given rigid symmetric monoidal ∞-category C. The resulting assignment from the Picard groupoid of C to ∞-categories with duality is lax symmetric monoidal (Corollary 8.5).

In section 9 we twist the duality further, this time by a C_2-action on the given rigid symmetric monoidal ∞-category C. Also in the case this C_2-action is trivial we get new twists, namely by C_2-automorphisms of the tensor unit of C (Remark 9.4). Our main result of this section is that hermitian K-theory twisted by a twisted second tensor power of a tensor invertible object and non-twisted hermitian K-theory coincide (Theorem 9.6 and Remark 9.15).

We combine K-theory and hermitian K-theory into a C_2-equivariant spectrum indexed on the trivial universe, see section 10.

We define the ∞-category of hermitian objects of an ∞-category with duality in section 11 generalizing the groupoidal versions we have used until this section in the paper.

In section 12 we discuss applications e.g. to finitely generated projective modules over E_∞-ring spectra making use of our twisting techniques to also define symplectic versions of hermitian K-theory.

In the paper [17] hermitian and real K-theory for stable ∞-categories with duality is developed using an ∞-categorical version of the hermitian $S_\bullet$-construction thereby obtaining a suitable theory which goes beyond direct sum hermitian K-theory. The method in loc. cit. to obtain a real K-theory spectrum can be equally well applied to the situation considered in this paper.

Expanded versions of parts of this paper will be part of the second named author's phd thesis.

Acknowledgements: We would like to thank Hongyi Chu, David Gepner, Jens Hornbostel, Kristian Moi, Thomas Nikolaus, Oliver Röndigs, Sean Tilson and Girja Tripathi for very helpful discussions and suggestions on the subject.

2 ∞-categorical background

Usually an ∞-category will mean a quasi category, i.e. a weak Kan complex. We frequently will have to change universes.

For $K, L \in \mathsf{sSet}$ we let $\mathrm{Fun}(K, L)$ denote L^K, especially when L is an ∞-category.

Spc denotes the ∞-category of spaces, i.e. $N(\mathsf{Kan})$, the nerve of the simplicial category of Kan complexes, see [10, Definition 1.2.16.1], and Sp the ∞-category

of spectra, i.e. the stabilzation of Spc. We denote by Cat_∞ the ∞-category of small ∞-categories, see [10, Definition 3.0.0.1], and by $\widehat{\mathsf{Cat}}_\infty$ the ∞-category of not necessarily small ∞-categories. We let Pr^L (resp. Pr^R) be the ∞-category of presentable ∞-categories and left (resp. right) adjoints as maps, see [10, Definition 5.5.3.1]. In particular $\mathsf{Spc}, \mathrm{Sp}, \mathsf{Cat}_\infty \in \mathsf{Pr}^L$.

For an ∞-category C and two objects $x, y \in C$ we let $\mathrm{Hom}_C(x,y)$ be a/the mapping space between x and y either considered as a Kan complex (with preferred model $C^{\Delta^1} \times_{C\times C} \{(x,y)\}$) or as an object in hoSpc.

A faithful functor $F\colon C \to D$ between ∞-categories is said to be good if for any $x, y \in C$ every component of $\mathrm{Hom}_D(F(x), F(y))$ on an equivalence is hit by an equivalence in $\mathrm{Hom}_C(x,y)$.

The good image of F is then defined to be the subcategory of D consisting of those vertices which are equivalent to vertices in the image of F and those edges that lie up to equivalence in the image of F.

The factorization of F through its good image is an equivalence followed by a subcategory.

A fully faithful functor is good, and in this case the essential image coincides with the good image.

Remark 2.1. *The good fully faithful functors are exactly the monomorphisms (in Cat_∞ or $\widehat{\mathsf{Cat}}_\infty$), where a map $X \to Y$ in an ∞-category is defined to be a monomorphism if the square*

$$\begin{array}{ccc} X & \longrightarrow & X \\ \downarrow & & \downarrow \\ X & \longrightarrow & Y \end{array}$$

is cartesian. Also the good fully faithful functors are exactly the faithful functors which induce fully faithful functors on core groupoids.

Proposition 2.2. *Let I be an ∞-category and let f be a monomorphism in Cat_∞^I. Then $\lim_I f$ is again a monomorphism whose good image can be characterized as follows: An object belongs to the good image if and only if its components lie in the good images of all the f_i, $i \in I$, and similarly for morphisms.*

Proposition 2.3. *Let $\varphi\colon C \to D$ be a symmetric monoidal functor between symmetric monoidal ∞-categories whose underlying functor is a good faithful functor. Then for any ∞-operad $\mathcal{O}$ (where for simplicity we assume that the fiber of $\mathcal{O}$ over $\langle 1\rangle$ is categorically contractible) the induced functor*

$$\mathrm{Alg}_{\mathcal{O}}(C) \to \mathrm{Alg}_{\mathcal{O}}(D)$$

is a good faithful functor. A morphism between $\mathcal{O}$-algebras belongs to its good image if and only if the underlying morphism belongs to the good image of φ.

Proposition 2.4. *Let $\varphi\colon C \to D$ be a good faithful functor between ∞-categories and let I be an ∞-category. Then the induced functor $\mathrm{Fun}(I,C) \to \mathrm{Fun}(I,D)$ is a good faithful functor whose good image can be characterized as follows: A functor $I \to D$ belongs to the good image if and only if it factors through the good image of φ, and similarly for morphisms.*

If C is a symmetric monoidal ∞-category and $A \in \mathrm{Alg}_{E_\infty}(C)$ then we let $\mathrm{Mod}_C(A)$ be the ∞-category of A-modules which we equip with its canonical symmetric monoidal structure if it exists. If it is clear from the context then we also write $\mathrm{Mod}(A)$ instead of $\mathrm{Mod}_C(A)$.

3 Infinity categories with duality

We begin with the

Theorem 3.1. *The (grouplike A_∞-) space of autoequivalences of Cat_∞ in Pr^L is (equivalent to) the discrete space $\mathbb{Z}/2$. The autoequivalence induced by the non-trivial element of $\mathbb{Z}/2$ sends a $C \in \mathsf{Cat}_\infty$ to C^{op}.*

Proof. This is [18, Th. 6.3] (see also [2, Theorem 8.12]). □

For a (discrete) group G and an ∞-category C we let $C[G]$ be the ∞-category of objects of C with G-action, i.e. the ∞-category $\mathrm{Fun}(BG, C)$. Here BG is the nerve of the category with one object and G as automorphisms of this object.

Pulling back along the map $BG \to *$ induces a map $C \to C[G]$ whose essential image is the full subcategory of objects with trivial G-action.

For $X \in C[G]$ we denote by X^{hG} the homotopy fixed points of the G-action on X, i.e. $\lim X \in C$ (where $X\colon BG \to C$ is the given functor), if this limit exists. If BG-shaped limits exist in C then $(-)^{hG}$ is right adjoint to $C \to C[G]$.

We let $C_2 := \mathbb{Z}/2$.

So Theorem 3.1 canonically exhibits Cat_∞ as an object of $\mathsf{Pr}^L[C_2]$.

Definition 3.2. *The ∞-category of small ∞-categories with duality is defined to be the homotopy fixed points of the C_2-action on Cat_∞, i.e. $\mathsf{Cat}_\infty^{hC_2} \in \mathsf{Pr}^L$.*

Theorem 3.3. *The C_2-action on Cat_∞ induces a C_2-action on the full subcategory $\mathsf{Spc} \subset \mathsf{Cat}_\infty$. This action is trivial.*

Proof. The maximal Kan subcomplex of $\mathrm{Fun}(\mathsf{Cat}_\infty, \mathsf{Cat}_\infty)$ on those vertices which are equivalences is a simplicial monoid M, and since C^{op} is an ∞-groupoid if $C \in \mathsf{Cat}_\infty$ is an ∞-groupoid M preserves $\mathsf{Spc} \subset \mathsf{Cat}_\infty$ by Theorem 3.1, establishing the first assertion.

The space of autoequivalences of Spc in Pr^L is contractible, establishing the second assertion. □

For C an ∞-category, $X \in C$ and $K \in \mathsf{Spc}$ we denote by $X^K \in C$ the cotensor if it exists. Clearly for $X \in C[G]$ (G a group) with trivial G-action we have $X^{hG} \simeq X^{BG}$.

It follows Spc^{hC_2} (with the trivial C_2-action from Theorem 3.3) is naturally equivalent to $\mathsf{Spc}[C_2]$, and we have canonical fully faithful maps $\mathsf{Spc}[C_2] \to \mathsf{Cat}_\infty^{hC_2}$ in Pr^R and Pr^L.

The fully faithfulness follows from the fact that the map $\mathsf{Spc} \to \mathsf{Cat}_\infty$ in $\mathsf{Pr}^L[C_2]$ has as a retraction in $\mathsf{Pr}^L[C_2]$ its left adjoint (or from the more general argument that a limit of fully faithful maps in $\widehat{\mathsf{Cat}}_\infty$ is again fully faithful).

Remark 3.4. *The full subcategory $\mathsf{Cat} \subset \mathsf{Cat}_\infty$ of ∞-categories which are equivalent to (small) categories is stable under the C_2-action giving a full embedding $\mathsf{Cat}^{hC_2} \to \mathsf{Cat}_\infty^{hC_2}$ in Pr^R.*

4 Symmetric monoidal structures

We will first show that the C_2-action on Cat_∞ is compatible with the natural symmetric monoidal structure on Cat_∞ (the cartesian product).

We let $\mathsf{SymMonCat}_\infty$ be the ∞-category of small symmetric monoidal ∞-categories and symmetric monoidal functors (one possible definition is as a subcategory of the ∞-category of ∞-operads, another one is as $\mathrm{Alg}_{E_\infty}(\mathsf{Cat}_\infty)$), and $\widehat{\mathsf{SymMonCat}}_\infty$ the ∞-category of not necessarily small symmetric monoidal ∞-categories.

We let $\mathsf{SymMonCat}_\infty^\times \subset \mathsf{SymMonCat}_\infty$ and $\widehat{\mathsf{SymMonCat}}_\infty^\times \subset \widehat{\mathsf{SymMonCat}}_\infty$ be the full subcategories of cartesian symmetric monoidal ∞-categories.

Let $\mathsf{Cat}_\infty^{\mathrm{Cart}}$ be the subcategory of Cat_∞ of small ∞-categories possessing finite products and product-preserving functors, and similarly for $\widehat{\mathsf{Cat}}_\infty^{\mathrm{Cart}}$.

Theorem 4.1. *The canonical forgetful functors*

$$\mathsf{SymMonCat}_\infty^\times \to \mathsf{Cat}_\infty^{\mathrm{Cart}}$$

and

$$\widehat{\mathsf{SymMonCat}}_\infty^\times \to \widehat{\mathsf{Cat}}_\infty^{\mathrm{Cart}}$$

are equivalences.

Proof. This is [9, Corollary 2.4.1.9]. □

Recall Pr^L has a natural symmetric monoidal structure such that an E_∞-algebra in Pr^L is a symmetric monoidal ∞-category whose underlying category is presentable and such that the tensor product preserves colimits separately in each variable. We denote the E_∞-algebras in Pr^L by $E_\infty(\mathsf{Pr}^L)$. Thus we have a natural good faithful functor

$$E_\infty(\mathsf{Pr}^L) \to \widehat{\mathsf{SymMonCat}}_\infty.$$

We let $E_\infty^\times(\mathsf{Pr}^L)$ be the full subcategory of $E_\infty(\mathsf{Pr}^L)$ on the cartesian symmetric monoidal presentable ∞-categories.

We get a commutative diagram

$$\begin{array}{ccccc} E_\infty(\mathsf{Pr}^L) & \longrightarrow & \widehat{\mathsf{SymMonCat}}_\infty & & \\ \uparrow & & \uparrow & & \\ E_\infty^\times(\mathsf{Pr}^L) & \xrightarrow{\varphi} & \widehat{\mathsf{SymMonCat}}_\infty^\times & \xrightarrow{\sim} & \widehat{\mathsf{Cat}}_\infty^{\mathrm{Cart}}. \end{array}$$

The ∞-category Cat_∞ together with its C_2-action can be viewed as a functor $BC_2 \to \widehat{\mathsf{Cat}}_\infty$ which factors through $\widehat{\mathsf{Cat}}_\infty^{\mathrm{Cart}}$.

Thus we naturally get (up to a contractible space of choices) a functor $\psi\colon BC_2 \to \widehat{\mathsf{SymMonCat}}_\infty^\times$ which lifts (up to a contractible space of choices) to a functor $BC_2 \to E_\infty^\times(\mathsf{Pr}^L)$, since ψ factors through the good image of φ. Its prolonguation

$$\psi'\colon BC_2 \to E_\infty(\mathsf{Pr}^L)$$

exhibits Cat_∞ as an object of $E_\infty(\mathsf{Pr}^L)[C_2]$.

Next we are going to consider bimonoidal structures with duality.

Recall from [4] the smashing localizations

$$C \mapsto C_*,\ C \mapsto \mathrm{Mon}_{E_\infty}(C),\ C \mapsto \mathrm{Grp}_{E_\infty}(C),\ C \mapsto \mathrm{Sp}(C)$$

of Pr^L.

Denoting by $\mathcal{L}$ any of these we get induced functors

$$E_\infty(\mathsf{Pr}^L) \xrightarrow{\mathcal{L}} E_\infty(\mathsf{Pr}^L)$$

which are localizations.

Precomposing with ψ' yields C_2-objects $\mathcal{L}(\mathsf{Cat}_\infty)$ in $E_\infty(\mathsf{Pr}^L)$, in particular

$$\mathsf{SymMonCat}_\infty \simeq \mathrm{Mon}_{E_\infty}(\mathsf{Cat}_\infty) \in E_\infty(\mathsf{Pr}^L)[C_2].$$

Remark 4.2. *As we did with* Cat_∞ *one can exhibit any cartesian closed presentable* ∞*-category with* C_2*-action as object of* $E_\infty(\mathsf{Pr}^L)[C_2]$*, in particular* Cat*. Applying* $\mathcal{L}$ *to* Cat *yields* $\mathsf{SymMonCat} \in E_\infty(\mathsf{Pr}^L)[C_2]$.

For $C \in E_\infty(\mathsf{Pr}^L)$ we let $E_\infty(C) := \mathrm{Alg}_{E_\infty}(C)$. The assignment $C \mapsto E_\infty(C)$ is a functor $e\colon E_\infty(\mathsf{Pr}^L) \to \mathsf{Pr}^L$. Precomposing this functor with the functor $BC_2 \to E_\infty(\mathsf{Pr}^L)$ corresponding to $\mathsf{SymMonCat}_\infty$ yields

$$\mathrm{Rig}_{E_\infty}(\mathsf{Cat}_\infty) := E_\infty(\mathsf{SymMonCat}_\infty) \in \mathsf{Pr}^L[C_2].$$

The ∞-category $E_\infty(\mathsf{Pr}^L)$ has (small) limits, thus we get an object

$$\mathsf{SymMonCat}_\infty^{hC_2} \in E_\infty(\mathsf{Pr}^L).$$

Proposition 4.3. *There is a natural equivalence*

$$E_\infty(\mathsf{SymMonCat}_\infty^{hC_2}) \simeq \mathrm{Rig}_{E_\infty}(\mathsf{Cat}_\infty)^{hC_2}$$

in Pr^L.

Proof. This follows from the fact that e preserves limits: Note first that it is sufficient to show that the functor $e'\colon E_\infty(\widehat{\mathsf{Cat}}_\infty) \to \widehat{\mathsf{Cat}}_\infty$, $C \mapsto E_\infty(C)$, preserves limits, since the inclusions $\mathsf{Pr}^L \to \widehat{\mathsf{Cat}}_\infty$ and $E_\infty(\mathsf{Pr}^L) \to E_\infty(\widehat{\mathsf{Cat}}_\infty)$ preserve limits.

The functor e' can be factored as

$$E_\infty(\widehat{\mathsf{Cat}}_\infty) \to \widehat{\mathrm{Op}}_\infty \to \widehat{\mathsf{Cat}}_\infty,$$

where $\widehat{\mathrm{Op}}_\infty$ is the ∞-category of not necessarily small ∞-operads, the first functor the natural inclusion and the second functor assigns to an ∞-operad $\mathcal{O}$ its ∞-category of commutative algebra objects in $\mathcal{O}$. But both functors have left adjoints, the first the symmetric monoidal envelope functor and the second the functor $C \mapsto C \otimes \mathrm{Comm}$, where $\otimes$ denotes the natural action of ∞-categories on ∞-operads and Comm the commutative operad. This shows e preserves limits. □

Remark 4.4. *Similarly to Proposition 4.3 we have an equivalence*

$$\mathsf{SymMonCat}_\infty^{hC_2} \simeq E_\infty(\mathsf{Cat}_\infty^{hC_2})$$

in Pr^L*, which can be promoted to an equivalence in* $E_\infty(\mathsf{Pr}^L)$*, since*

$$E_\infty(\mathsf{Cat}_\infty^{hC_2}) \simeq \mathrm{Mon}_{E_\infty}(\mathsf{Cat}_\infty^{hC_2})$$

($\mathsf{Cat}_\infty^{hC_2}$ *has the cartesian symmetric monoidal structure).*

The constructions done in this section starting from Cat_∞ can also be persued with Spc instead, and these resulting constructions are compatible with the map $\mathsf{Spc} \to \mathsf{Cat}_\infty$ in Pr^L.

In particular there is a map $\mathsf{Spc} \to \mathsf{Cat}_\infty$ in $E_\infty(\mathsf{Pr}^L)[C_2]$, yielding (by applying the smashing localization $\mathrm{Mon}_{E_\infty}(-)$) a map

$$\mathrm{Mon}_{E_\infty}(\mathsf{Spc}) \to \mathsf{SymMonCat}_\infty$$

in $E_\infty(\mathsf{Pr}^L)[C_2]$.

Taking E_∞-algebras yields a map

$$\mathrm{Rig}_{E_\infty}(\mathsf{Spc}) \coloneqq E_\infty(\mathrm{Mon}_{E_\infty}(\mathsf{Spc})) \to \mathrm{Rig}_{E_\infty}(\mathsf{Cat}_\infty)$$

in $\mathsf{Pr}^L[C_2]$, which finally gives a map

$$\mathrm{Rig}_{E_\infty}(\mathsf{Spc})^{hC_2} \to \mathrm{Rig}_{E_\infty}(\mathsf{Cat}_\infty)^{hC_2}$$

in Pr^L.

We note that there are equivalences

$$\mathrm{Rig}_{E_\infty}(\mathsf{Spc}[C_2]) \simeq E_\infty(\mathrm{Mon}_{E_\infty}(\mathsf{Spc})[C_2]) \simeq \mathrm{Rig}_{E_\infty}(\mathsf{Spc})[C_2] \simeq \mathrm{Rig}_{E_\infty}(\mathsf{Spc})^{hC_2}.$$

5 Group completions and spectra

We denote by

$$(-)^{\tilde{}} \colon \mathsf{Cat}_\infty \to \mathsf{Spc}$$

a right adjoint to the map $\mathsf{Spc} \to \mathsf{Cat}_\infty$ in Pr^L, and we denote by the same symbol similar right adjoints, e.g. the right adjoint to $\mathrm{Mon}_{E_\infty}(\mathsf{Spc}) \to \mathsf{SymMonCat}_\infty$.

The direct sum K-theory functor ([4, Definition 8.3]) is the composition

$$K \colon \mathsf{SymMonCat}_\infty \xrightarrow{(-)^{\tilde{}}} \mathrm{Mon}_{E_\infty}(\mathsf{Spc}) \to \mathrm{Grp}_{E_\infty}(\mathsf{Spc}) \to \mathrm{Sp},$$

which is lax symmetric monoidal [4, Theorem 8.6].

Its effect on rig-categories (E_∞-objects in $\mathsf{SymMonCat}_\infty$) can be described by the composition

$$\mathrm{Rig}_{E_\infty}(\mathsf{Cat}_\infty) \xrightarrow{(-)^{\tilde{}}} \mathrm{Rig}_{E_\infty}(\mathsf{Spc}) \to \mathrm{Ring}_{E_\infty}(\mathsf{Spc}) \to E_\infty(\mathrm{Sp}),$$

where $\mathrm{Ring}_{E_\infty}(\mathsf{Spc}) \coloneqq E_\infty(\mathrm{Grp}_{E_\infty}(\mathsf{Spc}))$.

We also denote by $(-)^{\tilde{}}$ a right adjoint of the map

$$\alpha \colon \mathrm{Mon}_{E_\infty}(\mathsf{Spc})[C_2] \simeq \mathrm{Mon}_{E_\infty}(\mathsf{Spc})^{hC_2} \to \mathsf{SymMonCat}_\infty^{hC_2}$$

in Pr^L.

We define the direct sum hermitian K-theory functor as the composition

$$K_h \colon \mathsf{SymMonCat}_\infty^{hC_2} \xrightarrow{(-)^{\tilde{}}} \mathrm{Mon}_{E_\infty}(\mathsf{Spc})[C_2] \xrightarrow{(-)^{hC_2}}$$
$$\mathrm{Mon}_{E_\infty}(\mathsf{Spc}) \to \mathrm{Grp}_{E_\infty}(\mathsf{Spc}) \to \mathrm{Sp}.$$

Proposition 5.1. *K_h is lax symmetric monoidal.*

Proof. Since α is in fact a map in $E_\infty(\mathsf{Pr}^L)$, the first map in the definition of K_h is lax symmetric monoidal. The second map is the right adjoint to the symmetric monoidal functor

$$\mathrm{Mon}_{E_\infty}(\mathsf{Spc}) \to \mathrm{Mon}_{E_\infty}(\mathsf{Spc})[C_2],$$

which sends an object to the object endowed with the trivial C_2-action. The third and fourth maps in the definition of K_h are symmetric monoidal. □

Note that the effect of K_h on E_∞-objects (rig-categories with duality) can be described by the composition

$$\mathrm{Rig}_{E_\infty}(\mathsf{Cat}_\infty)^{hC_2} \xrightarrow{(-)^\sim} \mathrm{Rig}_{E_\infty}(\mathsf{Spc})[C_2] \xrightarrow{(-)^{hC_2}}$$

$$\mathrm{Rig}_{E_\infty}(\mathsf{Spc}) \to \mathrm{Ring}_{E_\infty}(\mathsf{Spc}) \to E_\infty(\mathrm{Sp}).$$

6 The first hermitian recognition principle

We give a hermitian variant of [4, Theorem 8.8]. This section is heavily inspired by the proof of [4, Theorem 8.8].

We denote by $\mathsf{Cat}_\infty^\Sigma$ the subcategory of Cat_∞ of ∞-categories which admit finite coproducts and coproduct preserving functors. $\mathsf{Cat}_\infty^\Sigma$ is presentable and has a closed symmetric monoidal structure such that the left adjoint to the inclusion $\mathsf{Cat}_\infty^\Sigma \subset \mathsf{Cat}_\infty$ is symmetric monoidal (see the references in the proof of [4, Theorem 8.8] to [9]). $\mathsf{Cat}_\infty^\Sigma$ is preadditive (proof of [4, Theorem 8.8]), and the right adjoint to the composition

$$\beta\colon \mathsf{SymMonCat}_\infty \simeq \mathrm{Mon}_{E_\infty}(\mathsf{Cat}_\infty) \to \mathrm{Mon}_{E_\infty}(\mathsf{Cat}_\infty^\Sigma) \simeq \mathsf{Cat}_\infty^\Sigma$$

is the full embedding

$$\mathsf{Cat}_\infty^\Sigma \to \mathsf{SymMonCat}_\infty$$

which sends an ∞-category with coproducts to the corresponding cocartesian symmetric monoidal ∞-category (loc. cit.). Thus β is a symmetric monoidal localization.

The map β and its right adjoint are not compatible with the C_2-action on $\mathsf{SymMonCat}_\infty$, therefore we introduce another localization.

We let $i\colon \mathsf{Cat}_\infty^{\mathrm{preadd}} \subset \mathsf{Cat}_\infty^\Sigma$ be the full subcategory of preadditive categories, i.e. those categories which also have finite products, are pointed and in which for any two objects X and Y the natural map $X \coprod Y \to X \times Y$ is an equivalence.

Proposition 6.1. *The functor i has both a left and a right adjoint. The right adjoint is informally given by the assignment $C \mapsto \mathrm{Mon}_{E_\infty}(C^{\mathrm{op}})^{\mathrm{op}}$.*

Proof. We first note that the assignment $C \mapsto \mathrm{Mon}_{E_\infty}(C^{\mathrm{op}})^{\mathrm{op}}$ can be refined to a functor

$$r\colon \mathsf{Cat}_\infty^\Sigma \to \mathsf{Cat}_\infty^{\mathrm{preadd}}$$

(where we use in particular that $\mathrm{Mon}_{E_\infty}(C^{\mathrm{op}})$ is preadditive for $C \in \mathsf{Cat}_\infty^\Sigma$ by [4, Corollary 2.5. (i)]).

Moreover $r \circ i$ is naturally equivalent (say via φ) to the identity by [4, Proposition 2.3 (iv)]. For $C \in \mathsf{Cat}_\infty^\Sigma$ and $D \in \mathsf{Cat}_\infty^{\mathrm{preadd}}$ we have a chain of equivalences of ∞-categories

$$\mathrm{Fun}^{\amalg}(D, r(C)) \simeq \mathrm{Fun}^{\amalg}(D, \mathrm{Mon}_{E_\infty}(C^{\mathrm{op}})^{\mathrm{op}}) \simeq \mathrm{Fun}^{\prod}(D^{\mathrm{op}}, \mathrm{Mon}_{E_\infty}(C^{\mathrm{op}}))^{\mathrm{op}}$$

$$\simeq \mathrm{Fun}^{\prod}(D^{\mathrm{op}}, C^{\mathrm{op}})^{\mathrm{op}} \simeq \mathrm{Fun}^{\amalg}(D, C),$$

where the third equivalence follows from [4, Corollary 2.5 (iii)]. Here $\mathrm{Fun}^{\amalg}$ resp. $\mathrm{Fun}^{\prod}$ denotes finite coproducts resp. finite products preserving functors. It follows that φ exhibits r as a right adjoint to i.

Thus r is a colocalization, in particular the colocal objects in $\mathsf{Cat}_\infty^\Sigma$, i.e. $\mathsf{Cat}_\infty^{\mathrm{preadd}}$, are closed under colimits, thus $\mathsf{Cat}_\infty^{\mathrm{preadd}}$ has all small colimits and they are computed in $\mathsf{Cat}_\infty^\Sigma$.

Note that r preserves filtered colimits ($(-)^{\mathrm{op}}$ and $\mathrm{Mon}_{E_\infty}(-)$ preserve them). It follows that $\mathsf{Cat}_\infty^{\mathrm{preadd}}$ is presentable, thus, since i preserves colimits and limits (note that a limit of preadditive categories in $\mathsf{Cat}_\infty^\Sigma$ is preadditive), i also has a left adjoint. □

Let us denote a left adjoint of i by l.

Proposition 6.2. *The localization $i \circ l$ of $\mathsf{Cat}_\infty^\Sigma$ is symmetric monoidal.*

Proof. We have to check that the tensor product of a local equivalence f in $\mathsf{Cat}_\infty^\Sigma$ with an object $C \in \mathsf{Cat}_\infty^\Sigma$ is again a local equivalence. By mapping into local objects this is equivalent to the fact that for each preadditive $Z \in \mathsf{Cat}_\infty^\Sigma$ the internal Hom $\underline{\mathrm{Hom}}(C, Z)$ is preadditive, which can be verified directly. □

Thus $\mathsf{Cat}_\infty^{\mathrm{preadd}}$ is closed symmetric monoidal and l and $l \circ \beta$ are also symmetric monoidal localizations.

This time $l \circ \beta$ is compatible with the C_2-action on $\mathsf{SymMonCat}_\infty$ since its right adjoint

$$\mathsf{Cat}_\infty^{\mathrm{preadd}} \to \mathsf{SymMonCat}_\infty$$

is.

Theorem 6.3. *The functor*

$$l \circ \beta \colon \mathsf{SymMonCat}_\infty \to \mathsf{Cat}_\infty^{\mathrm{preadd}}$$

is naturally a map in $E_\infty(\mathsf{Pr}^L)[C_2]$.

Proof. Apply [9, Lemma 2.2.4.11] to the BC_2-cocartesian family

$$p \colon S^{\otimes} \to BC_2 \times \mathrm{N}\mathsf{Fin}_*$$

corresponding to the functor $BC_2 \to \widehat{\mathsf{SymMonCat}}_\infty$ encoding the C_2-action on the symmetric monoidal ∞-category $\mathsf{SymMonCat}_\infty$. □

Corollary 6.4. *There is a natural symmetric monoidal functor*

$$\mathsf{SymMonCat}_\infty^{hC_2} \to (\mathsf{Cat}_\infty^{\mathrm{preadd}})^{hC_2}$$

in Pr^L with lax symmetric monoidal right adjoint

$$(\mathsf{Cat}_\infty^{\mathrm{preadd}})^{hC_2} \to \mathsf{SymMonCat}_\infty^{hC_2}$$

in Pr^R.

We denote by $E_\infty^\Sigma(\mathsf{Cat}_\infty)$ the following subcategory of $E_\infty(\mathsf{Cat}_\infty)$: The objects are symmetric monoidal ∞-categories such that the underlying ∞-categories have finite coproducts and the given tensor products preserve these coproducts separately in each variable, and the maps are symmetric monoidal functors such that the underlying functors preserve finite coproducts.

We denote by $E_\infty^{\mathrm{preadd}}(\mathsf{Cat}_\infty) \subset E_\infty^\Sigma(\mathsf{Cat}_\infty)$ the full subcategory of those symmetric monoidal categories in $E_\infty^\Sigma(\mathsf{Cat}_\infty)$ whose underlying categories are preadditive.

Proposition 6.5. *The functor*

$$E_\infty(\mathsf{Cat}_\infty^\Sigma) \to E_\infty(\mathsf{Cat}_\infty)$$

factors through $E_\infty^\Sigma(\mathsf{Cat}_\infty)$*, the functor*

$$E_\infty(\mathsf{Cat}_\infty^{\mathrm{preadd}}) \to E_\infty(\mathsf{Cat}_\infty)$$

factors through $E_\infty^{\mathrm{preadd}}(\mathsf{Cat}_\infty)$*, and in the naturally induced commutative diagram*

$$\begin{array}{ccccc} E_\infty(\mathsf{Cat}_\infty) & \longleftarrow & E_\infty(\mathsf{Cat}_\infty^\Sigma) & \longleftarrow & E_\infty(\mathsf{Cat}_\infty^{\mathrm{preadd}}) \\ \downarrow{\scriptstyle =} & & \downarrow{\scriptstyle \sim} & & \downarrow{\scriptstyle \sim} \\ E_\infty(\mathsf{Cat}_\infty) & \longleftarrow & E_\infty^\Sigma(\mathsf{Cat}_\infty) & \longleftarrow & E_\infty^{\mathrm{preadd}}(\mathsf{Cat}_\infty) \end{array}$$

the vertical maps are equivalences.

Proof. The first statements are clear, the middle vertical equivalence is [9, Remark 4.8.1.9] and the right vertical equivalence follows from this equivalence. □

Proposition 6.6. *The functors*

$$E_\infty(\mathsf{Cat}_\infty) \to E_\infty(\mathsf{SymMonCat}_\infty) \to E_\infty(\mathsf{Cat}_\infty^{\mathrm{preadd}})$$

are naturally maps in $\mathsf{Pr}^L[C_2]$*, and its right adjoints*

$$E_\infty(\mathsf{Cat}_\infty^{\mathrm{preadd}}) \to E_\infty(\mathsf{SymMonCat}_\infty) \to E_\infty(\mathsf{Cat}_\infty)$$

are naturally maps in $\mathsf{Pr}^R[C_2]$*.*

Proof. The first statement follows from the fact that the functors

$$\mathsf{Cat}_\infty \to \mathsf{SymMonCat}_\infty \to \mathsf{Cat}_\infty^{\mathrm{preadd}}$$

are naturally maps in $E_\infty(\mathsf{Pr}^L)[C_2]$, and the second statement follows from the first. □

Proposition 6.7. *The* C_2*-action on* $E_\infty(\mathsf{Cat}_\infty)$ *induces a* C_2*-action on the* ∞*-category* $E_\infty^{\mathrm{preadd}}(\mathsf{Cat}_\infty)$*, and the induced functor*

$$E_\infty^{\mathrm{preadd}}(\mathsf{Cat}_\infty)^{hC_2} \to E_\infty(\mathsf{Cat}_\infty)^{hC_2}$$

is a good faithful functor whose good image consists of those objects and morphisms in $E_\infty(\mathsf{Cat}_\infty)^{hC_2}$ *whose underlying objects and morphisms in* $E_\infty(\mathsf{Cat}_\infty)$ *are in* $E_\infty^{\mathrm{preadd}}(\mathsf{Cat}_\infty)$*.*

Proposition 6.8. *The map*

$$E_\infty(\mathsf{Cat}_\infty^{\mathrm{preadd}}) \to E_\infty(\mathsf{Cat}_\infty)$$

in $\mathsf{Pr}^R[C_2]$ *factors via an equivalence*

$$E_\infty(\mathsf{Cat}_\infty^{\mathrm{preadd}}) \xrightarrow{\sim} E_\infty^{\mathrm{preadd}}(\mathsf{Cat}_\infty)$$

in $\mathsf{Pr}^R[C_2]$ *and* $\mathsf{Pr}^L[C_2]$.

Theorem 6.9. *Let* $C \in E_\infty(\mathsf{Cat}_\infty)^{hC_2} \simeq \mathsf{SymMonCat}_\infty^{hC_2}$*, such that the underlying* ∞*-category* C_u *is preadditive and such that the monoidal product*

$$\otimes\colon C_u \times C_u \to C_u$$

preserves finite coproducts separately in each variable. Then C *is canonically an object of* $\mathrm{Rig}_{E_\infty}(\mathsf{Cat}_\infty)^{hC_2}$*, i.e. a rig-category (or semiring* ∞*-category or bimonoidal* ∞*-category) with duality.*

Proof. By Proposition 6.7 C is naturally an object of $E_\infty^{\mathrm{preadd}}(\mathsf{Cat}_\infty)^{hC_2}$, so by Proposition 6.8 also an object of $E_\infty(\mathsf{Cat}_\infty^{\mathrm{preadd}})^{hC_2}$. Mapping C further along the functor

$$E_\infty(\mathsf{Cat}_\infty^{\mathrm{preadd}})^{hC_2} \to E_\infty(\mathsf{SymMonCat}_\infty)^{hC_2}$$

(see Proposition 6.6) in Pr^R yields the desired object. □

Combining Theorem 6.9 with the last composition in section 4 yields a map

$$E_\infty^{\mathrm{preadd}}(\mathsf{Cat}_\infty)^{hC_2} \to E_\infty(\mathrm{Sp}).$$

7 The second hermitian recognition principle

Let $\mathsf{SymMonCat}_\infty^{\mathrm{rig}} \subset \mathsf{SymMonCat}_\infty$ be the full subcategory of rigid symmetric monoidal ∞-categories. In the present section we will exhibit each object C of $\mathsf{SymMonCat}_\infty^{\mathrm{rig}}$ as a symmetric monoidal ∞-category with duality, where the duality is informally given by $C \ni X \mapsto X^\vee = \underline{\mathrm{Hom}}(X, \mathbf{1})$.

We need to use the theory of pairings from [9, Section 5.2.1]. But instead of right fibrations which appear there in the definition of pairing ([9, Definition 5.2.1.5]) we use left fibrations.

We let Pair be the full subcategory of $\mathrm{Fun}(\Lambda_0^2, \mathsf{Cat}_\infty)$ on those diagrams $C \leftarrow M \to D$ such that the induced map $M \to C \times D$ is equivalent to a left fibration. Pair is a localization of $\mathrm{Fun}(\Lambda_0^2, \mathsf{Cat}_\infty)$, see [9, Remark 5.2.1.15].

As in [9, Remark 5.2.1.20] there is the notion of perfect pairing, and we let $\mathsf{Pair}^{\mathrm{perf}} \subset \mathsf{Pair}$ be the subcategory of perfect pairings. The two forgetful functors $\phi, \psi\colon \mathsf{Pair}^{\mathrm{perf}} \to \mathsf{Cat}_\infty$ are equivalences, see [9, Remark 5.2.1.20].

The C_2-action on Λ_0^2 induces a strict C_2-action on Pair and on $\mathsf{Pair}^{\mathrm{perf}}$, and because of Theorem 3.1 ϕ and ψ commute with the C_2-actions.

Also the functor $(\phi, \psi)\colon \mathsf{Pair}^{\mathrm{perf}} \to \mathsf{Cat}_\infty \times \mathsf{Cat}_\infty$ strictly commutes with the C_2-actions, where we let C_2 act on $\mathsf{Cat}_\infty \times \mathsf{Cat}_\infty$ by permutation of the factors.

Lemma 7.1. *Let C be a small ∞-category and G a group acting on C. Then there is a natural equivalence*

$$(\mathsf{Cat}_\infty)^{\mathrm{lf}}_{/C} \simeq \mathrm{Fun}(C, \mathsf{Spc})$$

(where the superscript lf *denotes the full subcategory on those maps $D \to C$ which are equivalent to left fibrations) respecting the G-actions.*

For $C \in \mathsf{Cat}_\infty$ the fiber F_C of (ϕ, ψ) over (C, C) inherits a C_2-action. We note that F_C is an essentially small Kan complex.

Lemma 7.2. *For any $C \in \mathsf{Cat}_\infty$ there is a natural equivalence*

$$F_C \simeq (\mathsf{Cat}_\infty)^{\mathrm{perf}}_{/(C\times C)}$$

(where the superscript denotes the subcategory of perfect pairings and representable maps) respecting the C_2-actions.

Corollary 7.3. *To give a duality on $C \in \mathsf{Cat}_\infty$ is the same as to give a C_2-homotopy fixed point of* $\mathrm{Fun}(C\times C, \mathsf{Spc})$ *(or equivalently a functor $(C\times C)/^h C_2 \to \mathsf{Spc}$, where $/^h$ denotes homotopy orbits) which as a functor $C \times C \to \mathsf{Spc}$ is a perfect pairing.*

Proof. We show that the fiber F' of

$$\mathsf{Cat}_\infty^{hC_2} \to \mathsf{Cat}_\infty$$

over C is equivalent to $(\mathrm{Fun}(C \times C, \mathsf{Spc})^{\mathrm{perf}})^{hC_2}$.

Taking C_2-homotopy fixed points in the C_2-diagram

$$\begin{array}{ccc} F_C & \longrightarrow & \mathsf{Pair}^{\mathrm{perf}} \\ \downarrow & & \downarrow \\ * & \xrightarrow{(C,C)} & \mathsf{Cat}_\infty \times \mathsf{Cat}_\infty \end{array}$$

yields the (homotopy) pullback diagram

$$\begin{array}{ccc} F_C^{hC_2} & \longrightarrow & (\mathsf{Pair}^{\mathrm{perf}})^{hC_2} \\ \downarrow & & \downarrow \\ * & \xrightarrow{C} & \mathsf{Cat}_\infty, \end{array}$$

thus $F' \simeq F_C^{hC_2}$. The result follows now from Lemmas 7.1 and 7.2. □

For any $C \in \mathsf{Cat}_\infty$ we let $(\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/C}$ be the subcategory of $(\mathsf{Cat}_\infty)_{/C}$ on the maps $D \to C$ which are equivalent to cocartesian fibrations and the morphisms which are representable by cocartesian maps of cocartesian fibrations.

We let Pair_C be the full subcategory of $\mathrm{Fun}(\Lambda^2_0, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/C})$ on those functors whose corresponding functors $C \to \mathrm{Fun}(\Lambda^2_0, \mathsf{Cat}_\infty)$ factor through Pair. We denote by $\mathsf{Pair}_C^{\mathrm{perf}}$ the subcategory of Pair_C on the functors which correspond to functors $C \to \mathsf{Pair}$ which factor through $\mathsf{Pair}^{\mathrm{perf}}$ and the morphisms whose

corresponding natural transformations between functors $C \to \mathsf{Pair}$ also factor through $\mathsf{Pair}^{\mathrm{perf}}$.

Note Pair_C and $\mathsf{Pair}_C^{\mathrm{perf}}$ inherit natural C_2-actions.

We let Fin_* be the standard skeleton of the category of finite pointed sets.

For each $C \in \mathsf{SymMonCat}_\infty^{\mathrm{rig}}$ we will produce an object in $(\mathsf{Pair}^{\mathrm{perf}}_{\mathrm{N}\mathsf{Fin}_*})^{hC_2}$ whose corresponding functor

$$\mathrm{N}\mathsf{Fin}_* \to (\mathsf{Pair}^{\mathrm{perf}})^{hC_2}$$

exhibits C as a symmetric monoidal ∞-category with duality.

We let $\mathcal{C}_1$ be the subcategory of Fin_* generated by the objects $\langle 1\rangle$ and $\langle 2\rangle$ and the following morphisms: all automorphisms of $\langle 2\rangle$, the unique active map $\langle 2\rangle \to \langle 1\rangle$ and all inert maps $\langle 2\rangle \to \langle 1\rangle$.

We let $\mathcal{C}_1'$ be the same subcategory of Fin_* as $\mathcal{C}_1$ except that we remove the active morphism $\langle 2\rangle \to \langle 1\rangle$.

We let $\mathcal{C}_2$ be the category arising by glueing $\mathcal{C}_1'$ to $[1]$ along the objects $\langle 2\rangle$ and 0 subject to the relation that there results a unique map from $\langle 2\rangle = 0$ to 1.

We extend the inclusion $\mathcal{C}_1' \to \mathcal{C}_1$ to a functor $f\colon \mathcal{C}_2 \to \mathcal{C}_1$ by sending $1 \in [1]$ to $\langle 1\rangle$ and the unique non-identity map in $[1]$ to the active morphism $\langle 2\rangle \to \langle 1\rangle$.

Let $\mathcal{C}_3$ be the category arising by glueing $\mathcal{C}_2$ to $[1]$ along the objects $1 \in [1] \subset \mathcal{C}_2$ and $1 \in [1]$.

We let ι be the composition $\mathcal{C}_1 \to \mathsf{Fin}_* \hookrightarrow \mathsf{Spc}_*$.

For any $C \in E_\infty(\mathsf{Cat}_\infty)$ we obtain a functor $\iota_C\colon \mathcal{C}_1 \to E_\infty(\mathsf{Cat}_\infty)$ defined by $x \mapsto \iota(x) \wedge C$, where $\wedge$ denotes the tensoring of pointed spaces on $E_\infty(\mathsf{Cat}_\infty)$ (note that $E_\infty(\mathsf{Cat}_\infty)$ is pointed).

There is a natural fully faithful functor $j'\colon E_\infty(\mathsf{Cat}_\infty) \to (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathrm{N}\mathsf{Fin}_*}$ whose essential image we denote by $(\mathsf{Cat}_\infty)^{\mathrm{sm}}_{/\mathrm{N}\mathsf{Fin}_*}$. We denote the induced equivalence

$$E_\infty(\mathsf{Cat}_\infty) \to (\mathsf{Cat}_\infty)^{\mathrm{sm}}_{/\mathrm{N}\mathsf{Fin}_*}$$

by j.

There is a natural forgetful functor $\nu\colon (\mathsf{Cat}_\infty)^{\mathrm{sm}}_{/\mathrm{N}\mathsf{Fin}_*} \to \mathsf{Cat}_{\infty,*}$ to pointed ∞-categories sending a map $C^\otimes \to \mathrm{N}\mathsf{Fin}_*$ to $C^\otimes$, where the pointing is given by the homotopically unique object in the fiber over $\langle 0\rangle$ (which "corresponds" to the tensor unit).

Let $\mathsf{Cat}_\infty^{\Delta^1,\mathrm{lf}}$ be the full subcategory of $\mathsf{Cat}_\infty^{\Delta^1}$ of maps which are equivalent to left fibrations.

We let $\mu\colon \mathsf{Cat}_{\infty,*} \to \mathsf{Cat}_\infty^{\Delta^1,\mathrm{lf}}$ which sends (C, X) to the left fibration corresponding to the functor $C \ni Y \mapsto \mathrm{Hom}(X, Y)$.

In the following when we apply functor categories we sometimes leave out taking the nerve.

We let

$$k_1\colon \mathrm{Fun}(\mathcal{C}_2, (\mathsf{Cat}_\infty)^{\mathrm{sm}}_{/\mathrm{N}\mathsf{Fin}_*}) \times_{\mathsf{Cat}_\infty} \mathsf{Cat}_\infty^{\Delta^1,\mathrm{lf}} \to \mathrm{Fun}(\mathcal{C}_3, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathrm{N}\mathsf{Fin}_*}),$$

where the fiber product is via evaluation at the object a of $\mathcal{C}_2$ which does not belong to $\mathcal{C}_1'$ and evaluation at the target of an object in $\mathsf{Cat}_\infty^{\Delta^1,\mathrm{lf}}$, be the functor which "glues" the given map in $\mathsf{Cat}_\infty^{\Delta^1,\mathrm{lf}}$ via its target to a.

Precomposing k_1 with the functor

$$\mathrm{Fun}(\mathcal{C}_2, (\mathsf{Cat}_\infty)^{\mathrm{sm}}_{/\mathrm{N}\mathsf{Fin}_*}) \to \mathrm{Fun}(\mathcal{C}_2, (\mathsf{Cat}_\infty)^{\mathrm{sm}}_{/\mathrm{N}\mathsf{Fin}_*}) \times_{\mathsf{Cat}_\infty} \mathsf{Cat}_{\infty,*}$$

$$\to \mathrm{Fun}(\mathcal{C}_2, (\mathsf{Cat}_\infty)^{\mathrm{sm}}_{/\mathsf{NFin}_*}) \times_{\mathsf{Cat}_\infty} \mathsf{Cat}_\infty^{\Delta^1,\mathrm{lf}},$$

where the first functor is the identity in the first component and evaluation at a followed by ν in the second component, and the second functor the identity in the first component and μ in the second, yields a functor

$$k_2 \colon \mathrm{Fun}(\mathcal{C}_2, (\mathsf{Cat}_\infty)^{\mathrm{sm}}_{/\mathsf{NFin}_*}) \to \mathrm{Fun}(\mathcal{C}_3, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*}).$$

Sending a $C \in E_\infty(\mathsf{Cat}_\infty)$ to $j \circ \iota_C \circ f$ yields a functor

$$\gamma \colon E_\infty(\mathsf{Cat}_\infty) \to \mathrm{Fun}(\mathcal{C}_2, (\mathsf{Cat}_\infty)^{\mathrm{sm}}_{/\mathsf{NFin}_*}),$$

composing this with k_2 yields

$$k_3 \colon E_\infty(\mathsf{Cat}_\infty) \to \mathrm{Fun}(\mathcal{C}_3, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*}).$$

We let Ψ' be the category arising by glueing Λ^2_0 to $[1]$ along the objects $0 \in \Lambda^2_0$ and $0 \in [1]$. Also let Ψ be the category arising by glueing Ψ' to $[1]$ along the objects $1 \in [1] \subset \Psi'$ and $1 \in [1]$.

Ψ' and Ψ have C_2-actions, where C_2 acts on Λ^2_0 as above and on the remaining parts of the categories trivially.

Let I be the category with two objects 0 and 1 and a unique isomorphism from 0 to 1. C_2 acts on I by permuting 0 and 1.

We let C_2 act on $I \times \Psi$ diagonally.

There is a natural functor $(I \times \Psi)/C_2 \to \mathcal{C}_3$, where the quotient in the source is in the 1-categorical sense, described as follows:

On $(I \times \Lambda^2_0)/C_2$ it factors through $\mathcal{C}'_1$ and is then an equivalence, it sends the image in $(I \times \Psi')/C_2$ of the objects in $I \times \Psi'$ not belonging to $I \times \Lambda^2_0$ to a, and the remaining object in $(I \times \Psi)/C_2$ to the remaining object in $\mathcal{C}_3$.

Thus we obtain a C_2-equivariant functor $I \times \Psi \to \mathcal{C}_3$ for the trivial C_2-action on $\mathcal{C}_3$.

This induces a functor

$$\mathrm{Fun}(\mathcal{C}_3, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*}) \to \mathrm{Fun}(I \times \Psi, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})^{hC_2}.$$

The projection $I \times \Psi \to \Psi$ is a C_2-equivariant underlying equivalence, we thus have an equivalence

$$\mathrm{Fun}(\Psi, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})^{hC_2} \simeq \mathrm{Fun}(I \times \Psi, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})^{hC_2}.$$

Choosing an inverse of this equivalence yields

$$\mathrm{Fun}(\mathcal{C}_3, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*}) \to \mathrm{Fun}(\Psi, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})^{hC_2},$$

and composing with k_3 yields

$$k_4 \colon E_\infty(\mathsf{Cat}_\infty) \to \mathrm{Fun}(\Psi, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})^{hC_2}.$$

We let $\widetilde{\Psi}$ be the category obtained from Ψ by glueing in an additional Λ^2_0, where the "endpoints" of Λ^2_0 are glued to the objects $0 \in \Lambda^2_0 \subset \Psi$ and the object in Ψ which is not in Ψ' and where the resulting square should commute by definition (this glueing produces additional morphisms, for example there are also unique morphisms form 0 in the new Λ^2_0 to the endpoints of the old Λ^2_0).

There is an induced C_2-action on $\widetilde{\Psi}$ which fixes everything new.

We let $\square \subset \widetilde{\Psi}$ be the square formed by the new Λ^2_0 and those (non-identity) morphisms of Ψ which are not in the old Λ^2_0. Let

$$k_5 \colon \mathrm{Fun}^{\square}(\widetilde{\Psi}, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*}) \subset \mathrm{Fun}(\widetilde{\Psi}, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})$$

be the full subcategory on those diagrams which become cartesian after restricting to $\square \subset \widetilde{\Psi}$, then the restriction functor

$$\mathrm{Fun}^{\square}(\widetilde{\Psi}, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*}) \to \mathrm{Fun}(\Psi, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})$$

is an equivalence. Choosing an inverse and composing with k_5 yields a C_2-map

$$\mathrm{Fun}(\Psi, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*}) \to \mathrm{Fun}(\widetilde{\Psi}, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})$$

inducing a map

$$\mathrm{Fun}(\Psi, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})^{hC_2} \to \mathrm{Fun}(\widetilde{\Psi}, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})^{hC_2}$$

whose composition with k_4 yields

$$k_6 \colon E_\infty(\mathsf{Cat}_\infty) \to \mathrm{Fun}(\widetilde{\Psi}, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})^{hC_2}.$$

Let $g \colon \Lambda^2_0 \to \widetilde{\Psi}$ be the C_2-equivariant functor which sends $0 \in \Lambda^2_0$ to 0 in the new Λ^2_0 in $\widetilde{\Psi}$ and the two endpoints of Λ^2_0 to the two endpoints of the old Λ^2_0 in $\widetilde{\Psi}$.

Restriction along g produces from k_6 a map

$$k_7 \colon E_\infty(\mathsf{Cat}_\infty) \to \mathrm{Fun}(\Lambda^2_0, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})^{hC_2}.$$

Remark 7.4. *For $C \in E_\infty(\mathsf{Cat}_\infty)$ and $\langle n \rangle \in \mathsf{Fin}_*$, forgetting homotopy fixed points and taking the fiber over $\langle n \rangle$ yields from $k_7(C)$ a diagram $\Lambda^2_0 \to \mathsf{Cat}_\infty$ which has the shape*

$$C^n \leftarrow M \to C^n$$

such that the induced map $M \xrightarrow{\varphi} C^n \times C^n$ is (equivalent to) a left fibration and the fiber of φ over an object $((X_1, \dots, X_n), (Y_1, \dots, Y_n))$ is equivalent to

$$\mathrm{Hom}_C(\mathbf{1}, X_1 \otimes Y_1) \times \cdots \times \mathrm{Hom}_C(\mathbf{1}, X_n \otimes Y_n).$$

Lemma 7.5. *The restriction of k_7 to $\mathsf{SymMonCat}^{\mathrm{rig}}_\infty$ factors through the good image of the good faithful functor*

$$(\mathsf{Pair}^{\mathrm{perf}}_{\mathsf{NFin}_*})^{hC_2} \to \mathrm{Fun}(\Lambda^2_0, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})^{hC_2},$$

giving rise to a functor

$$k_8 \colon \mathsf{SymMonCat}^{\mathrm{rig}}_\infty \to (\mathsf{Pair}^{\mathrm{perf}}_{\mathsf{NFin}_*})^{hC_2} \simeq \mathrm{Fun}(\mathsf{NFin}_*, \mathsf{Pair}^{\mathrm{perf}})^{hC_2}$$

$$\simeq \mathrm{Fun}(\mathsf{NFin}_*, (\mathsf{Pair}^{\mathrm{perf}})^{hC_2}) \simeq \mathrm{Fun}(\mathsf{NFin}_*, \mathsf{Cat}^{hC_2}_\infty).$$

Proof. This follows by construction (see also Remark 7.4). □

Lemma 7.6. *The functor*

$$\mathsf{SymMonCat}_\infty^{\mathrm{rig}} \to \mathrm{Fun}(\mathrm{N}\mathsf{Fin}_*, \mathsf{Cat}_\infty)$$

induced by k_8 by forgetting homotopy fixed points in the target is naturally equivalent to the composition

$$\mathsf{SymMonCat}_\infty^{\mathrm{rig}} \subset \mathsf{SymMonCat}_\infty \to \mathrm{Fun}(\mathrm{N}\mathsf{Fin}_*, \mathsf{Cat}_\infty),$$

where the second functor is the standard one.

Proof. This follows by construction. □

We have proved:

Theorem 7.7. *The forgetful functor*

$$(\mathsf{SymMonCat}_\infty^{\mathrm{rig}})^{hC_2} \to \mathsf{SymMonCat}_\infty^{\mathrm{rig}}$$

has a natural section.

Combining with Theorem 6.9 yields

Theorem 7.8. *Let C be a rigid symmetric monoidal ∞-category, such that the underlying ∞-category C_u is preadditive and such that the monoidal product*

$$\otimes : C_u \times C_u \to C_u$$

preserves finite coproducts separately in each variable. Then C is canonically an object of $\mathrm{Rig}_{E_\infty}(\mathsf{Cat}_\infty)^{hC_2}$, i.e. a rig-category (or semiring ∞-category or bimonoidal ∞-category) with duality.

Let $\mathsf{SymMonCat}_\infty^{\mathrm{rig,preadd}}$ be the full subcategory of $E_\infty^{\mathrm{preadd}}(\mathsf{Cat}_\infty)$ of rigid symmetric monoidal ∞-categories.

Corollary 7.9. *We have a natural hermitian K-theory functor*

$$\mathsf{SymMonCat}_\infty^{\mathrm{rig,preadd}} \to E_\infty(\mathrm{Sp}).$$

8 The extended second hermitian recognition principle

This section is a variation of the previous section. We will equip any rigid symmetric monoidal ∞-category C with a duality of the form $X \mapsto X^\vee \otimes L$, where L is a tensor invertible object of C.

We keep the notation of the last section.

For an ∞-category C we let α_C be the composition

$$(\mathsf{Cat}_\infty)_{/C}^{\mathrm{cocart}} \to \mathrm{Fun}(C, \mathsf{Cat}_\infty) \overset{(\psi^{-1})_*}{\longrightarrow} \mathrm{Fun}(C, \mathsf{Pair}^{\mathrm{perf}}) \to \mathrm{Fun}(C, \mathsf{Pair})$$

$$\to \mathrm{Fun}(C, \mathrm{Fun}(\Lambda_0^2, \mathsf{Cat}_\infty)) \simeq \mathrm{Fun}(C \times \Lambda_0^2, \mathsf{Cat}_\infty) \simeq \mathrm{Fun}(\Lambda_0^2, (\mathsf{Cat}_\infty)_{/C}^{\mathrm{cocart}}).$$

α_C followed by evaluation at $2 \in \Lambda_0^2$ is naturally equivalent to the identity.

We let $\mathcal{C}_4$ be the category obtained by glueing $\mathcal{C}_2$ to Λ^2_0 along the object of $\mathcal{C}_2$ which does not belong to $\mathcal{C}'_1$ and the object $2 \in \Lambda^2_0$.

We have a natural composition

$$\beta_C : \mathrm{Fun}(\mathcal{C}_2, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/C})$$
$$\to \mathrm{Fun}(\mathcal{C}_2, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/C}) \times_{(\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/C}} \mathrm{Fun}(\Lambda^2_0, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/C})$$
$$\simeq \mathrm{Fun}(\mathcal{C}_4, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/C}),$$

where the first functor is the identity into the first factor and the functor α_C into the second.

Thus we get an induced functor

$$h_1 : E_\infty(\mathsf{Cat}_\infty) \xrightarrow{\gamma} \mathrm{Fun}(\mathcal{C}_2, (\mathsf{Cat}_\infty)^{\mathrm{sm}}_{/\mathsf{NFin}_*})$$
$$\to \mathrm{Fun}(\mathcal{C}_2, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*}) \xrightarrow{\beta_{\mathsf{NFin}_*}} \mathrm{Fun}(\mathcal{C}_4, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*}).$$

Let Θ be the category obtained by glueing Ψ' and Λ^2_0 along the objects $1 \in [1] \subset \Psi'$ and $2 \in \Lambda^2_0$. The group C_2 acts on Θ by its action on Ψ' and the trivial action on Λ^2_0. There is a natural C_2-equivariant full embedding $\Psi \to \Theta$.

There is a natural functor $(I \times \Theta)/C_2 \to \mathcal{C}_4$ extending the natural functor $(I \times \Psi')/C_2 \to \mathcal{C}_2$ from the previous section.

As in the previous section we thus obtain

$$\delta : \mathrm{Fun}(\mathcal{C}_4, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*}) \to \mathrm{Fun}(\Theta, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})^{hC_2},$$

and precomposing with h_1 yields

$$h_2 : E_\infty(\mathsf{Cat}_\infty) \to \mathrm{Fun}(\Theta, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})^{hC_2}.$$

Let x be the object of $\widetilde{\Psi}$ which lies in Ψ but not in Ψ', and let $\widetilde{\Theta}$ be the category obtained by glueing $\widetilde{\Psi}$ and $[1]$ along the objects x and 0.

There is a canonical C_2-equivariant full embedding $\Theta \to \widetilde{\Theta}$ extending the embedding $\Psi \to \widetilde{\Psi}$.

We let

$$\mathrm{Fun}^{\square}(\widetilde{\Theta}, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*}) \subset \mathrm{Fun}(\widetilde{\Theta}, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})$$

be the full subcategory such that the restriction of the given diagram to the square of $\widetilde{\Theta}$ is cartesian.

The restriction functor

$$\mathrm{Fun}^{\square}(\widetilde{\Theta}, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*}) \to \mathrm{Fun}(\Theta, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})$$

is an equivalence, thus, choosing an inverse, yields the C_2-functors

$$\mathrm{Fun}(\Theta, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*}) \to \mathrm{Fun}^{\square}(\widetilde{\Theta}, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})$$
$$\to \mathrm{Fun}(\widetilde{\Theta}, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})$$

and the induced functor

$$\mathrm{Fun}(\Theta, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})^{hC_2} \to \mathrm{Fun}(\widetilde{\Theta}, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})^{hC_2}.$$

Precomposing with h_2 yields

$$h_3 \colon E_\infty(\mathsf{Cat}_\infty) \to \mathrm{Fun}(\widetilde{\Theta}, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})^{hC_2}.$$

We let $\tilde{g} \colon \Psi' \to \widetilde{\Theta}$ be the functor which extends g and sends $1 \in [1] \subset \Psi'$ to the object of $\widetilde{\Theta}$ which does not lie in $\widetilde{\Psi}$.

Restriction along $\tilde{g}$ and precomposing with h_3 yields

$$h_4 \colon E_\infty(\mathsf{Cat}_\infty) \to \mathrm{Fun}(\Psi', (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})^{hC_2}.$$

Remark 8.1. *For $C \in E_\infty(\mathsf{Cat}_\infty)$ and $\langle n \rangle \in \mathsf{Fin}_*$, forgetting homotopy fixed points and taking the fiber over $\langle n \rangle$ yields from $h_4(C)$ a diagram $\Psi' \to \mathsf{Cat}_\infty$ which has the shape*

$$\begin{array}{ccccc} & & (C^{\mathrm{op}})^n & & \\ & & \uparrow & & \\ C^n & \longleftarrow & M & \longrightarrow & C^n \end{array}$$

such that the induced map $M \xrightarrow{\varphi} C^n \times C^n \times (C^{\mathrm{op}})^n$ is (equivalent to) a left fibration and the fiber of φ over an object $((X_1, \dots, X_n), (Y_1, \dots, Y_n), (Z_1, \dots, Z_n))$ is equivalent to

$$\mathrm{Hom}_C(Z_1, X_1 \otimes Y_1) \times \cdots \times \mathrm{Hom}_C(Z_n, X_n \otimes Y_n).$$

Let P be the category obtained by glueing two copies of $[2]$ along the morphisms $0 \to 1$ in both of these categories. Let us call m the morphism in P which is the morphism $0 \to 2$ in the first copy of $[2]$ and m' the one in the second.

Let $p \colon \Lambda^2_0 \to P$ be the functor which send $0 \in \Lambda^2_0$ to the two glued 1's and the endpoints of Λ^2_0 to the two 2's.

Let Q be the category obtained by glueing two copies of P along the morphisms m in both of the copies. The image of the two m's in Q is again denoted by m. The morphisms m' in the two copies of P give morphisms m_1 and m_2 in Q.

Permuting the two copies of P yields a C_2-action on Q.

The functor p gives rise to two functors $p_1, p_2 \colon \Lambda^2_0 \to Q$.

Let $\Psi' \to Q$ be the C_2-functor which sends the two legs of $\Lambda^2_0 \subset \Psi'$ to m_1 and m_2 and the morphism in $[1] \subset \Psi'$ to m.

Let

$$\mathrm{Fun}^{\times}(Q, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*}) \subset \mathrm{Fun}(Q, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})$$

be the full subcategory of such diagrams whose restrictions to Λ^2_0 via the functors p_1 and p_2 are product diagrams.

Then the restriction

$$\mathrm{Fun}^{\times}(Q, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*}) \to \mathrm{Fun}(\Psi', (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})$$

is an equivalence. Choosing an inverse yields the C_2-functor

$$\mathrm{Fun}(\Psi', (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*}) \to \mathrm{Fun}(Q, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathsf{NFin}_*})$$

which induces

$$\mathrm{Fun}(\Psi', (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathrm{N}\mathsf{Fin}_*})^{hC_2} \to \mathrm{Fun}(Q, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathrm{N}\mathsf{Fin}_*})^{hC_2}.$$

Precomposing with h_4 gives

$$h_5 \colon E_\infty(\mathsf{Cat}_\infty) \to \mathrm{Fun}(Q, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathrm{N}\mathsf{Fin}_*})^{hC_2}.$$

Let $R \subset Q$ be the full subcategory which contains m and the two possible factorizations of m in Q. R inherits a C_2 action.

The functor h_5 induces by restriction to R finally a functor

$$h_6 \colon E_\infty(\mathsf{Cat}_\infty) \to \mathrm{Fun}(R, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathrm{N}\mathsf{Fin}_*})^{hC_2}.$$

Note R is C_2-isomorphic to $[1] \times [1]$, where C_2 acts on $[1] \times [1]$ by permutation.

Restricting to $(1,1)$ via this isomorphism and precomposing with h_6 sends C to C^{op} with its opposite symmetric monoidal structure.

Note restriction of an $h_6(C)$ to $(1,1)$ yields in fact an object of

$$(\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathrm{N}\mathsf{Fin}_*}[C_2],$$

but by construction the C_2-action is trivial (this follows already from the definition of δ).

The functor h_6 induces the composition

$$\begin{aligned} v_1 \colon * &\to \mathrm{Fun}(E_\infty(\mathsf{Cat}_\infty), \mathrm{Fun}(R, (\mathsf{Cat}_\infty)^{\mathrm{cocart}}_{/\mathrm{N}\mathsf{Fin}_*})^{hC_2}) \\ &\simeq \mathrm{Fun}(E_\infty(\mathsf{Cat}_\infty), \mathrm{Fun}(R, \mathrm{Fun}(\mathrm{N}\mathsf{Fin}_*, \mathsf{Cat}_\infty))^{hC_2}) \\ &\simeq \mathrm{Fun}(E_\infty(\mathsf{Cat}_\infty) \times R \times \mathrm{N}\mathsf{Fin}_*, \mathsf{Cat}_\infty)^{hC_2} \\ &\simeq \mathrm{Fun}(R, \mathrm{Fun}(E_\infty(\mathsf{Cat}_\infty) \times \mathrm{N}\mathsf{Fin}_*, \mathsf{Cat}_\infty))^{hC_2} \\ &\to \mathrm{Fun}(R, (\widehat{\mathsf{Cat}}_\infty)^{\mathrm{cocart}}_{/(E_\infty(\mathsf{Cat}_\infty) \times \mathrm{N}\mathsf{Fin}_*)})^{hC_2}. \end{aligned}$$

Composing further via the functor induced by the forgetful functor

$$(\widehat{\mathsf{Cat}}_\infty)^{\mathrm{cocart}}_{/(E_\infty(\mathsf{Cat}_\infty) \times \mathrm{N}\mathsf{Fin}_*)} \to \widehat{\mathsf{Cat}}_\infty$$

yields

$$v_2 \colon * \to \mathrm{Fun}(R, \widehat{\mathsf{Cat}}_\infty)^{hC_2}.$$

Let $\mathrm{Fun}(R, \widehat{\mathsf{Cat}}_\infty)^{\mathrm{cocart}} \subset \mathrm{Fun}(R, \widehat{\mathsf{Cat}}_\infty)$ be the full subcategory on those diagrams d where $d(\varphi)$ is equivalent to a cocartesian fibration for any map φ in R whose target is final.

Then v_2 factors as a map

$$v_3 \colon * \to (\mathrm{Fun}(R, \widehat{\mathsf{Cat}}_\infty)^{\mathrm{cocart}})^{hC_2},$$

and the evaluation X at $(1,1)$ of v_3 has trivial C_2-action.

Lemma 8.2. *Let G be a group acting on an ∞-category C and let D be the ∞-category obtained from C by adjoining a final object. D carries an induced G-action. Let*

$$\mathrm{Fun}(D, \widehat{\mathsf{Cat}}_\infty)^{\mathrm{cocart}} \subset \mathrm{Fun}(D, \widehat{\mathsf{Cat}}_\infty)$$

be the full subcategory on those diagrams d where $d(\varphi)$ is equivalent to a cocartesian fibration for any map φ in D whose target is final.

Let $q \in \widehat{\mathsf{Cat}}_\infty$ and F be the fiber over q of the functor

$$\mathrm{Fun}(D, \widehat{\mathsf{Cat}}_\infty)^{\mathrm{cocart}} \to \widehat{\mathsf{Cat}}_\infty$$

which is evaluation at the final object. F inherits a G-action.

Then there is a natural G-equivariant equivalence

$$F \simeq \mathrm{Fun}(C, (\widehat{\mathsf{Cat}}_\infty)^{\mathrm{cocart}'}_{/q}),$$

where $(\widehat{\mathsf{Cat}}_\infty)^{\mathrm{cocart}'}_{/q} \subset (\widehat{\mathsf{Cat}}_\infty)_{/q}$ is the full subcategory on maps $p \to q$ which are equivalent to cocartesian fibrations.

We thus get a cartesian C_2-equivariant square

$$\begin{array}{ccc} \mathrm{Fun}(\Lambda^2_0, (\widehat{\mathsf{Cat}}_\infty)^{\mathrm{cocart}'}_{/X}) & \longrightarrow & \mathrm{Fun}(R, \widehat{\mathsf{Cat}}_\infty)^{\mathrm{cocart}} \\ \downarrow & & \downarrow \\ \{X\} & \longrightarrow & \widehat{\mathsf{Cat}}_\infty \end{array}$$

(here $\widehat{\mathsf{Cat}}_\infty$ carries the trivial C_2-action), where the bottom C_2-map is obtained from a map with trivial C_2-action.

Taking homotopy fixed points yields the cartesian square

$$\begin{array}{ccc} \mathrm{Fun}(\Lambda^2_0, (\widehat{\mathsf{Cat}}_\infty)^{\mathrm{cocart}'}_{/X})^{hC_2} & \longrightarrow & (\mathrm{Fun}(R, \widehat{\mathsf{Cat}}_\infty)^{\mathrm{cocart}})^{hC_2} \\ \downarrow & & \downarrow \\ \{X\} & \longrightarrow & \widehat{\mathsf{Cat}}_\infty[C_2], \end{array}$$

so v_3 induces a functor

$$v_4 \colon * \to \mathrm{Fun}(\Lambda^2_0, (\widehat{\mathsf{Cat}}_\infty)^{\mathrm{cocart}'}_{/X})^{hC_2}.$$

This functor factors in fact via a functor

$$v_5 \colon * \to \mathrm{Fun}(\Lambda^2_0, (\widehat{\mathsf{Cat}}_\infty)^{\mathrm{cocart}}_{/X})^{hC_2}.$$

But we have

$$\mathrm{Fun}(\Lambda^2_0, (\widehat{\mathsf{Cat}}_\infty)^{\mathrm{cocart}}_{/X})^{hC_2} \simeq \mathrm{Fun}(\Lambda^2_0 \times X, \widehat{\mathsf{Cat}}_\infty)^{hC_2} \simeq \mathrm{Fun}(X, \mathrm{Fun}(\Lambda^2_0, \widehat{\mathsf{Cat}}_\infty)^{hC_2}),$$

and the prolonguation of v_5 along these equivalences gives a factorization

$$v_6 \colon * \to \mathrm{Fun}(X, \mathsf{Pair}^{hC_2}).$$

We let

$$\widetilde{\mathcal{U}} \to E_\infty(\mathsf{Cat}_\infty) \times \mathsf{NFin}_*$$

be the opposite universal fibration, i.e. the cocartesian fibration classified by the functor

$$c\colon E_\infty(\mathsf{Cat}_\infty)\times \mathrm{N}\mathsf{Fin}_* \to \mathrm{Fun}(\mathrm{N}\mathsf{Fin}_*, \mathsf{Cat}_\infty)\times \mathrm{N}\mathsf{Fin}_* \to \mathsf{Cat}_\infty \to \mathsf{Cat}_\infty,$$

where the first functor sends a symmetric monoidal ∞-category to the corresponding functor $\mathrm{N}\mathsf{Fin}_* \to \mathsf{Cat}_\infty$ in the first component and is the identity in the second, the scond functor is evaluation and the third one is $(-)^{\mathrm{op}}$.

Note that evaluation of v_1 at $(1,1)$ is equivalent to $\widetilde{\mathcal{U}}$ over $E_\infty(\mathsf{Cat}_\infty)\times \mathrm{N}\mathsf{Fin}_*$ and that thus $X \simeq \widetilde{\mathcal{U}}$ as ∞-categories.

Let $\widetilde{\mathcal{U}}^{\mathrm{rig}}$ be the pullback of $\widetilde{\mathcal{U}}$ along the functor

$$\mathsf{SymMonCat}_\infty^{\mathrm{rig}} \times \mathrm{N}\mathsf{Fin}_* \to E_\infty(\mathsf{Cat}_\infty)\times \mathrm{N}\mathsf{Fin}_*.$$

We let $\mathrm{Pic}(\widetilde{\mathcal{U}}^{\mathrm{rig}}) \to \widetilde{\mathcal{U}}^{\mathrm{rig}}$ be the family of Picard groupoids of $\widetilde{\mathcal{U}}^{\mathrm{rig}}$.

Lemma 8.3. *The composition*

$$\mathrm{Pic}(\widetilde{\mathcal{U}}^{\mathrm{rig}}) \to \widetilde{\mathcal{U}}^{\mathrm{rig}} \to \widetilde{\mathcal{U}} \simeq X \xrightarrow{v_6} \mathsf{Pair}^{hC_2}$$

factors through $(\mathsf{Pair}^{\mathrm{perf}})^{hC_2}$.

Proof. This follows by construction: Let $C \in \mathsf{SymMonCat}_\infty^{\mathrm{rig}}$, $\langle n\rangle \in \mathsf{Fin}_*$, then the fiber of

$$\mathrm{Pic}(\widetilde{\mathcal{U}}^{\mathrm{rig}}) \to \mathsf{SymMonCat}_\infty^{\mathrm{rig}} \times \mathrm{N}\mathsf{Fin}_*$$

over $(C, \langle n\rangle)$ is equivalent to $(\mathrm{Pic}(C)^{\mathrm{op}})^n$, where $\mathrm{Pic}(C)$ denotes the Picard groupoid of C. Let $(L_1, \dots, L_n) \in (\mathrm{Pic}(C)^{\mathrm{op}})^n$, so we get an object $\mathcal{L} = (C, \langle n\rangle, (L_1, \dots, L_n)) \in \mathrm{Pic}(\widetilde{\mathcal{U}}^{\mathrm{rig}})$.

Forgetting homotopy fixed points, the image of $\mathcal{L}$ under the composition in the Lemma has the shape

$$C^n \leftarrow M \to C^n$$

such that the induced map $M \xrightarrow{\varphi} C^n \times C^n$ is (equivalent to) a left fibration and the fiber of φ over an object $((X_1, \dots, X_n), (Y_1, \dots, Y_n))$ is equivalent to

$$\mathrm{Hom}_C(L_1, X_1 \otimes Y_1) \times \cdots \times \mathrm{Hom}_C(L_n, X_n \otimes Y_n).$$

Since all the L_i are $\otimes$-invertible the claim follows. □

We note that Picard groupoids are equivalent to its opposites, thus $\mathrm{Pic}(\widetilde{\mathcal{U}}^{\mathrm{rig}})$ is equivalent to the Picard groupoid of the pullback of the universal fibration over $E_\infty(\mathsf{Cat}_\infty)\times \mathrm{N}\mathsf{Fin}_*$ to $\mathsf{SymMonCat}_\infty^{\mathrm{rig}} \times \mathrm{N}\mathsf{Fin}_*$ which we denote by $\mathrm{Pic}(\mathcal{U}^{\mathrm{rig}})$.

We thus get by Lemma 8.3 an induced functor

$$v_7\colon \mathrm{Pic}(\mathcal{U}^{\mathrm{rig}}) \to \mathsf{Cat}_\infty^{hC_2}.$$

Lemma 8.4. *The functor v_7 defines a family of lax symmetric monoidal functors to $\mathsf{Cat}_\infty^{hC_2}$ endowed with the cartesian symmetric monoidal structure.*

Proof. This follows from the fact that the composition of v_7 to Cat_∞ is equivalent to the functor

$$\mathrm{Pic}(\mathcal{U}^{\mathrm{rig}}) \to \mathsf{SymMonCat}_\infty^{\mathrm{rig}} \times \mathrm{N}\mathsf{Fin}_* \to E_\infty(\mathsf{Cat}_\infty)\times \mathrm{N}\mathsf{Fin}_* \to \mathsf{Cat}_\infty,$$

where the last functor is c except that we do not take $(-)^{\mathrm{op}}$ in the end. □

Corollary 8.5. *i) Let $C \in \mathsf{SymMonCat}_\infty^{\mathrm{rig}}$ and denote by $\mathrm{Pic}(C)$ the Picard groupoid of C. Then there is an induced lax symmetric monoidal functor*

$$T_C\colon \mathrm{Pic}(C) \to \mathsf{Cat}_\infty^{hC_2}$$

whose composition to Cat_∞ is the constant functor at C.

In particular the image of the tensor unit defines an object of $\mathsf{SymMonCat}^{hC_2}$ which is the same as the object defined in the previous section, $T_C(L)$ is a module over $T_C(\mathbf{1})$ for any $L \in \mathrm{Pic}(C)$ and for $L, L' \in \mathrm{Pic}(C)$ there are natural multiplication maps $T_C(L) \times T_C(L') \to T_C(L \otimes L')$ which are commutative and associative. For $L \in \mathrm{Pic}(C)$ the duality on C is informally given by $C \ni X \mapsto X^\vee \otimes L$.

ii) For a map $C \to D$ in $\mathsf{SymMonCat}_\infty^{\mathrm{rig}}$ there is a natural transformation between lax symmetric monoidal functors from T_C to the composition $\mathrm{Pic}(C) \to \mathrm{Pic}(D) \xrightarrow{T_D} \mathsf{Cat}^{hC_2}$.

Remark 8.6. *Let $\mathrm{Pic}(\mathcal{U}^{\mathrm{rig}})_1$ be the restriction to $\langle 1 \rangle$ of the natural projection $\mathrm{Pic}(\mathcal{U}^{\mathrm{rig}}) \to \mathsf{NFin}_*$. Then $\mathrm{Pic}(\mathcal{U}^{\mathrm{rig}})_1$ can be viewed as the category with objects (C, L), where $C \in \mathsf{SymMonCat}_\infty^{\mathrm{rig}}$ and $L \in \mathrm{Pic}(C)$. The restriction v_8 of v_7 to $\mathrm{Pic}(\mathcal{U}^{\mathrm{rig}})_1$ preserves products.*

So only the functor v_8 is necessary to deduce the lax symmetric monoidal functors T_C:

The map $\pi\colon \mathrm{Pic}(\mathcal{U}^{\mathrm{rig}})_1 \to \mathsf{SymMonCat}_\infty^{\mathrm{rig}}$ is equivalent to a cocartesian (even left) fibration and preserves products. It is then possible to equip the fibers of π with symmetric monoidal structures (which are the same as the given ones) such that the inclusions of the fibers into $\mathrm{Pic}(\mathcal{U}^{\mathrm{rig}})_1$ are lax symmetric monoidal. Composition with v_8 is then also lax symmetric monoidal.

Proposition 8.7. *Let C be an ∞-category and $\mathcal{P} \to C \times \mathsf{NFin}_*$ a cocartesian family of ∞-operads. The good faithful lax symmetric monoidal functor*

$$(\mathsf{Cat}_\infty^{\mathrm{preadd}})^{hC_2} \to \mathsf{Cat}_\infty^{hC_2}$$

induces a good faithful functor

$$\mathrm{Alg}_{\mathcal{P}}((\mathsf{Cat}_\infty^{\mathrm{preadd}})^{hC_2}) \to \mathrm{Alg}_{\mathcal{P}}(\mathsf{Cat}_\infty^{hC_2}),$$

whose good image consists of those algebras A such that for any $c \in C$ the induced lax symmetric monoidal functor $A_c\colon \mathcal{P}_c \to \mathsf{Cat}_\infty$ satisfies the following properties: the underlying functor of A_c factors through $\mathsf{Cat}_\infty^{\mathrm{preadd}}$, and for any $x, y \in \mathcal{P}_c$ the functor

$$A_c(x) \times A_c(y) \to A_c(x \otimes y)$$

preserves finite coproducts separately in each variable, and those maps whose components in Cat_∞ preserve finite coproducts.

Let $\mathrm{Pic}(\mathcal{U}^{\mathrm{rig,preadd}})$ be the pullback of $\mathrm{Pic}(\mathcal{U}^{\mathrm{rig}})$ along the functor

$$\mathsf{SymMonCat}_\infty^{\mathrm{rig,preadd}} \times \mathsf{NFin}_* \to \mathsf{SymMonCat}_\infty^{\mathrm{rig}} \times \mathsf{NFin}_*.$$

Composition with v_7 exhibits an object of $\mathrm{Alg}_{\mathrm{Pic}(\mathcal{U}^{\mathrm{rig,preadd}})}(\mathsf{Cat}_\infty^{hC_2})$, and we deduce from Proposition 8.7 that this comes from an object of

$$\mathrm{Alg}_{\mathrm{Pic}(\mathcal{U}^{\mathrm{rig,preadd}})}((\mathsf{Cat}_\infty^{\mathrm{preadd}})^{hC_2}).$$

Corollary 8.8. *For any $C \in \mathsf{SymMonCat}_\infty^{\mathrm{rig,preadd}}$ the functor T_C factors via a lax symmetric monoidal functor $\mathrm{Pic}(C) \to (\mathsf{Cat}_\infty^{\mathrm{preadd}})^{hC_2}$. The composition*

$$T_C^{\mathrm{Sp}}\colon \mathrm{Pic}(C) \to (\mathsf{Cat}_\infty^{\mathrm{preadd}})^{hC_2} \to \mathsf{SymMonCat}_\infty^{hC_2} \xrightarrow{K_h} \mathrm{Sp},$$

where the second map is from Corollary 6.4, consists of lax symmetric monoidal functors.

In particular the image of the tensor unit is an E_∞-ring spectrum which is the same as the corresponding object in the previous section, $T_C^{\mathrm{Sp}}(L)$ is a module over $T_C^{\mathrm{Sp}}(\mathbf{1})$ for any $L \in \mathrm{Pic}(C)$ and for $L, L' \in \mathrm{Pic}(C)$ there are natural multiplication maps $T_C^{\mathrm{Sp}}(L) \otimes T_C^{\mathrm{Sp}}(L') \to T_C^{\mathrm{Sp}}(L \otimes L')$ which are commutative and associative.

Naturality with respect to maps in $\mathsf{SymMonCat}_\infty^{\mathrm{rig,preadd}}$ is given.

9 Twisting the duality

We keep the notation of the last section.

Let C be an ∞-category with an action of a group G.

If G acts on an object of C^{hG} one can obtain a twisted homotopy fixed point of C by considering the composition

$$t_C\colon C^{hG}[G] \simeq (C^{hG})^{hG} \simeq C^{h(G\times G)} \to C^{hG},$$

where $G \times G$ acts on C via the first projection and where the last arrow is induced by restriction to G via the diagonal $G \to G \times G$ (note that the composition of this diagonal followed by the first projection is the identity of G, thus the induced action of G on C is the original one).

Remark 9.1. *Let $G = C_2$ and $\tau\colon C \to C$ the action of the nontrivial element of C_2. Let $X \in C^{hC_2}[C_2]$ and denote by X also the underlying object in C. Since X is in particular a homotopy fixed point for the action of C_2 on C we have an equivalence*

$$f\colon X \to \tau(X)$$

in C such that $\tau(f) \circ f\colon X \to \tau^2(X) \simeq X$ is naturally equivalent to the identity in C. The additional C_2-action on $X \in C^{hC_2}$ gives us an equivalence $g\colon X \to X$ in C with natural equivalences $f \circ g \simeq \tau(g) \circ f$ and $f^2 \simeq \mathrm{id}$.

The underlying equivalence $X \to \tau(X)$ in C of the new fixed point datum on $t_C(X)$ is then $g \circ f$.

Specializing to the case $C = \mathsf{Cat}_\infty$, $G = C_2$ and the canonical action of C_2 on Cat_∞ we obtain the functor

$$t\colon \mathsf{Cat}_\infty^{hC_2}[C_2] \to \mathsf{Cat}_\infty^{hC_2}$$

and to the case $C = \mathsf{SymMonCat}_\infty$ the functor

$$t^{\mathrm{sm}}\colon \mathsf{SymMonCat}_\infty^{hC_2}[C_2] \to \mathsf{SymMonCat}_\infty^{hC_2}.$$

Pulling back the functor

$$\mathrm{Pic}(\mathcal{U}^{\mathrm{rig}})[C_2] \to (\mathsf{SymMonCat}_\infty^{\mathrm{rig}} \times \mathsf{NFin}_*)[C_2] \simeq \mathsf{SymMonCat}_\infty^{\mathrm{rig}}[C_2] \times \mathrm{NFin}_*[C_2]$$

to $\mathsf{SymMonCat}_\infty^{\mathrm{rig}}[C_2]\times\mathrm{N}\mathsf{Fin}_*$ yields the family of (twisted) Picard groupoids

$$F^t\colon \mathrm{Pic}(\mathcal{U}^{\mathrm{rig}})^t \to \mathsf{SymMonCat}_\infty^{\mathrm{rig}}[C_2]\times\mathrm{N}\mathsf{Fin}_*.$$

Note that the fiber of F_1^t (the restriction of F^t to $\mathsf{SymMonCat}_\infty^{\mathrm{rig}}[C_2]\times\{\langle 1\rangle\}$) over a rigid symmetric monoidal ∞-category C with C_2-action is $\mathrm{Pic}(C)^{hC_2}$ (where we use the induced C_2-action on $\mathrm{Pic}(C)$).

The composition

$$\mathrm{Pic}(\mathcal{U}^{\mathrm{rig}})^t \to \mathrm{Pic}(\mathcal{U}^{\mathrm{rig}})[C_2] \xrightarrow{v_7[C_2]} \mathsf{Cat}_\infty^{hC_2}[C_2]$$

followed by t gives us a functor

$$v_9\colon \mathrm{Pic}(\mathcal{U}^{\mathrm{rig}})^t \to \mathsf{Cat}_\infty^{hC_2}$$

defining, as in Lemma 8.4, a family of lax symmetric monoidal functors to $\mathsf{Cat}_\infty^{hC_2}$. So for any $C\in\mathsf{SymMonCat}_\infty^{\mathrm{rig}}[C_2]$ we obtain a lax symmetric monoidal functor

$$T_C\colon \mathrm{Pic}(C)^{hC_2}\to\mathsf{Cat}_\infty^{hC_2}.$$

Remark 9.2. *In this situation let $L\in\mathrm{Pic}(C)^{hC_2}$ and $f\colon C\to C$ the equivalence of C in $\mathsf{SymMonCat}_\infty^{\mathrm{rig}}$ induced by the nontrivial element of C_2. Then the duality on C underlying $T_C(L)$ is informally given by $C\ni X\mapsto f(X)^\vee\otimes L$.*

Pulling back $\mathrm{Pic}(\mathcal{U}^{\mathrm{rig}})^t$ further to $\mathsf{SymMonCat}_\infty^{\mathrm{rig,preadd}}[C_2]\times\mathrm{N}\mathsf{Fin}_*$ defines the family

$$\mathrm{Pic}(\mathcal{U}^{\mathrm{rig,preadd}})^t \to \mathsf{SymMonCat}_\infty^{\mathrm{rig,preadd}}[C_2]\times\mathrm{N}\mathsf{Fin}_*,$$

and composition with v_9 exhibits an object of

$$\mathrm{Alg}_{\mathrm{Pic}(\mathcal{U}^{\mathrm{rig,preadd}})^t}(\mathsf{Cat}_\infty^{hC_2})$$

which comes, using Proposition 8.7, from an object of

$$\mathrm{Alg}_{\mathrm{Pic}(\mathcal{U}^{\mathrm{rig,preadd}})^t}((\mathsf{Cat}_\infty^{\mathrm{preadd}})^{hC_2}).$$

Corollary 9.3. *For any $C\in\mathsf{SymMonCat}_\infty^{\mathrm{rig,preadd}}[C_2]$ we obtain a composition*

$$T_C^{\mathrm{Sp}}\colon \mathrm{Pic}(C)^{hC_2}\to(\mathsf{Cat}_\infty^{\mathrm{preadd}})^{hC_2}\to\mathsf{SymMonCat}_\infty^{hC_2}\xrightarrow{K_h}\mathrm{Sp}$$

which consists of lax symmetric monoidal functors.

Remark 9.4. *Let $C\in\mathsf{SymMonCat}_\infty^{\mathrm{rig}}$ and equip it with the trivial C_2-action. Then $\mathrm{Pic}(C)^{hC_2}\simeq\mathrm{Pic}(C)[C_2]\simeq\mathrm{Pic}(C)\times A$, where A is the fiber of the map $\mathrm{Pic}(C)[C_2]\to\mathrm{Pic}(C)$ which forgets the C_2-action. Thus we can twist the duality on the object of $(\mathsf{SymMonCat}_\infty^{\mathrm{rig}})^{hC_2}$ which, via Theorem 7.7, C gives rise to, by objects of A. Note that the underlying space of A is the space of A_∞-maps from C_2 to the space of (homotopy) automorphisms of the tensor unit of $\mathrm{Pic}(C)$.*

We are now interested in relations among the different types of duality twists we have encountered so far (i.e. by tensor invertible objects and C_2-automorphisms of the tensor unit).

To indicate what will be a consequence of the main result let $L \in \mathrm{Pic}(C)$, where $C \in \mathsf{SymMonCat}_\infty^{\mathrm{rig}}$. Then $L^{\otimes 2}$ is naturally an element of $\mathrm{Pic}(C)[C_2]$, where the C_2-action ι on $L \otimes L$ switches the two factors.

We will show that then $T_C((L^{\otimes 2}, \iota))$ and $T_C(\mathbf{1})$ are naturally equivalent in $\mathsf{Cat}_\infty^{hC_2}$.

The C_2-action on $L^{\otimes 2}$ can be defined using the following general construction:

Let D be an ∞-category which is preadditive and complete.

Let $X \in D$. For any natural number n we will factor the multiplication map $n\colon X \to X$ in a canonical way through $X^{B\Sigma_n}$.

Lemma 9.5. *Let the notation be as above. Let $\underline{n} := \{1, \dots, n\}$. Then there is a natural equivalence*

$$\underline{n} \cdot X \simeq X^{\underline{n}}$$

in $D[\Sigma_n]$, where the Σ_n-actions are induced via the canonical action of Σ_n on $\underline{n}$.

When we view X to be equiped with the trivial Σ_n-action the natural maps $X \to X^{\underline{n}}$ and $\underline{n} \cdot X \to X$ induced by the projection of $\underline{n}$ to the point are Σ_n-equivariant.

Thus we obtain the composition

$$m_X(n)\colon X \to (X^{\underline{n}})^{h\Sigma_n} \simeq (\underline{n} \cdot X)^{h\Sigma_n} \to X^{h\Sigma_n} \simeq X^{B\Sigma_n}$$

in D, and if we compose this map with the natural map $X^{B\Sigma_n} \to X$ this is the multiplication by n map.

If we let $D = \mathsf{SymMonCat}_\infty$ then for any $C \in \mathsf{SymMonCat}_\infty$ we obtain the symmetric monoidal functor

$$m_C(n)\colon C \to C[\Sigma_n]$$

which sends any $X \in C$ to $X^{\otimes n}$ with the natural Σ_n-action and which is natural in C.

For $n = 2$ and $L \in \mathrm{Pic}(C)$ we obtain the object $(L^{\otimes 2}, \iota)$ from above.

Now let $C \in \mathsf{SymMonCat}_\infty^{\mathrm{rig}}[C_2]$, then we obtain the symmetric monoidal functor

$$m_{\mathrm{Pic}(C)^{hC_2}}(2)\colon \mathrm{Pic}(C)^{hC_2} \to \mathrm{Pic}(C)^{hC_2}[C_2],$$

and composing with the (symmetric monoidal) functor

$$t_{\mathrm{Pic}(C)}\colon \mathrm{Pic}(C)^{hC_2}[C_2] \to \mathrm{Pic}(C)^{hC_2}$$

yields a symmetric monoidal functor

$$sq_C\colon \mathrm{Pic}(C)^{hC_2} \to \mathrm{Pic}(C)^{hC_2}$$

sending an L to $(L^{\otimes 2})'$, where the prime indicates that the C_2-homotopy fixed point data is twisted by the action of C_2 on $L^{\otimes 2}$.

Classical periodicity theorems for Witt- and Grothendieck-Witt groups are predecessors of the following

Theorem 9.6. *Let $C \in \mathsf{SymMonCat}_\infty^{\mathrm{rig}}[C_2]$ and $L \in \mathrm{Pic}(C)$. Then $T_C(sq_C(L))$ and $T_C(\mathbf{1})$ are canonically equivalent in $\mathsf{Cat}_\infty^{hC_2}$.*

To prepare for the proof we start with the following

Remark 9.7. *Let G be a group and C an ∞-category. By adjunction to give an object in $C[G]$, i.e. a map $* \to C[G]$ in $\widehat{\mathsf{Cat}}_\infty$, is the same as to give a map $* \to C$ in $\widehat{\mathsf{Cat}}_\infty[G]$, where we equip C with the trivial G-action.*

Now let $C \in \mathsf{SymMonCat}_\infty^{\mathrm{rig}}[C_2]$ and view it as a map $* \to \mathsf{SymMonCat}_\infty^{\mathrm{rig}}[C_2]$ in $\widehat{\mathsf{Cat}}_\infty$. This induces a diagram

$$\begin{array}{ccc} & & \mathrm{Pic}(\mathcal{U}^{\mathrm{rig}})[C_2] \\ & & \downarrow \\ * \times \mathrm{NFin}_* & \longrightarrow & \mathsf{SymMonCat}_\infty^{\mathrm{rig}}[C_2] \times \mathrm{NFin}_*[C_2] \end{array}$$

in $\widehat{\mathsf{Cat}}_\infty$, which via Remark 9.7 induces the diagram

$$\begin{array}{ccc} & & \mathrm{Pic}(\mathcal{U}^{\mathrm{rig}}) \\ & & \downarrow \\ * \times \mathrm{NFin}_* & \longrightarrow & \mathsf{SymMonCat}_\infty^{\mathrm{rig}} \times \mathrm{NFin}_* \end{array}$$

in $\widehat{\mathsf{Cat}}_\infty[C_2]$.

Taking the pullback yields

$$\begin{array}{ccc} \mathrm{Pic}(C)^{\otimes} & \longrightarrow & \mathrm{Pic}(\mathcal{U}^{\mathrm{rig}}) \\ \downarrow & & \downarrow \\ * \times \mathrm{NFin}_* & \longrightarrow & \mathsf{SymMonCat}_\infty^{\mathrm{rig}} \times \mathrm{NFin}_* \end{array}$$

in $\widehat{\mathsf{Cat}}_\infty[C_2]$.

Prolonging with the map $\mathrm{Pic}(\mathcal{U}^{\mathrm{rig}}) \to \mathsf{Cat}_\infty^{hC_2}$ yields a map

$$\mu_C \colon \mathrm{Pic}(C) \to \mathsf{Cat}_\infty^{hC_2}$$

which is a lax symmetric monoidal functor with C_2-action (where the action on the target is trivial).

Taking C_2-homotopy fixed points gives

$$\mathrm{Pic}(C)^{hC_2} \to \mathsf{Cat}_\infty^{hC_2}[C_2],$$

and prolonging with t gives T_C.

Proposition 9.8. *Let G be a group and $D \in E_\infty(\widehat{\mathsf{Cat}}_\infty)[G]$. Let $A \in E_\infty(D)^{hG} \simeq E_\infty(D^{hG})$. Then there is a natural G-action on the symmetric monoidal ∞-category $\mathrm{Mod}_D(A)$ and an equivalence*

$$\mathrm{Mod}_{D^{hG}}(A) \simeq \mathrm{Mod}_D(A)^{hG}$$

of symmetric monoidal ∞-categories.

Proof. Let $\mathcal{O}$ be an ∞-operad such that an algebra over $\mathcal{O}$ in a symmetric monoidal ∞-category C consists of an E_∞-algebra B in C together with a B-module M.

The E_∞-algebra A corresponds to a map $* \to \mathrm{Alg}_{E_\infty}(D)$ in $\widehat{\mathsf{Cat}}_\infty[G]$, so we get a pullback diagram

$$\begin{array}{ccc} \mathrm{Mod}_D(A) & \longrightarrow & \mathrm{Alg}_{\mathcal{O}}(D) \\ \downarrow & & \downarrow \\ * & \longrightarrow & \mathrm{Alg}_{E_\infty}(D) \end{array}$$

in $\widehat{\mathsf{Cat}}_\infty[G]$.

Taking G-homotopy fixed points yields the diagram

$$\begin{array}{ccc} \mathrm{Mod}_D(A)^{hG} & \longrightarrow & \mathrm{Alg}_{\mathcal{O}}(D^{hG}) \\ \downarrow & & \downarrow \\ * & \longrightarrow & \mathrm{Alg}_{E_\infty}(D^{hG}) \end{array}$$

in $\widehat{\mathsf{Cat}}_\infty$, inducing an equivalence

$$\mathrm{Mod}_D(A)^{hG} \simeq \mathrm{Mod}_{D^{hG}}(A).$$

The symmetric monoidality of this equivalence follows by fibering everything over $\mathrm{N}\mathsf{Fin}_*$. □

Let again $C \in \mathsf{SymMonCat}_\infty^{\mathrm{rig}}[C_2]$. Then by the second hermitian recognition principle we obtain an object

$$C \in \mathsf{SymMonCat}_\infty^{hC_2}[C_2] \simeq E_\infty(\mathsf{Cat}_\infty^{h(C_2\times C_2)}).$$

Forgetting the additional C_2-action we have an object $C' \in \mathsf{SymMonCat}_\infty^{hC_2}$, and it follows from Proposition 9.8 that there is a natural C_2-action on

$$\mathrm{Mod}_{\mathsf{Cat}_\infty^{hC_2}}(C') \simeq \mathrm{Mod}_{\mathsf{Cat}_\infty}(C)^{hC_2^1}$$

(since $C \in E_\infty(\mathsf{Cat}_\infty^{h(C_2\times C_2)})$, Proposition 9.8 yields an action of $C_2 \times C_2$ on $\mathrm{Mod}_{\mathsf{Cat}_\infty}(C)$, and C_2^1 refers to this action restricted to the first factor).

Now the lax symmetric monoidal C_2-functor μ_C can be factored as lax symmetric monoidal C_2-functors

$$\mathrm{Pic}(C) \xrightarrow{\tilde{\mu}_C} \mathrm{Mod}_{\mathsf{Cat}_\infty^{hC_2}}(C') \to \mathsf{Cat}_\infty^{hC_2},$$

where the second functor forgets the C'-module structure.

Lemma 9.9. *The functor $\tilde{\mu}_C$ is symmetric monoidal, so it can be considered as a map in $\widehat{\mathsf{SymMonCat}}_\infty[C_2]$.*

Proof. This is detected by mapping further to $\mathrm{Mod}_{\mathsf{Cat}_\infty}(C)$ and noting that all objects of $\mathrm{Pic}(C)$ are mapped after this prolonguation to the tensor unit in $\mathrm{Mod}_{\mathsf{Cat}_\infty}(C)$ (we also use that the funcor $\mathsf{Cat}_\infty^{hC_2} \to \mathsf{Cat}_\infty$ is conservative). □

We let C_2 act on $\mathrm{Mod}_{\mathsf{Cat}_\infty}(C)$ by restricting the $C_2 \times C_2$-action to the diagonal of $C_2 \times C_2$.

By taking C_2-homotopy fixed points the functor $\tilde{\mu}_C$ induces a functor

$$\widetilde{T}_C \colon \mathrm{Pic}(C)^{hC_2} \to \mathrm{Mod}_{\mathsf{Cat}_\infty}(C)^{h(C_2 \times C_2)} \to \mathrm{Mod}_{\mathsf{Cat}_\infty}(C)^{hC_2},$$

where the second functor is restriction along the diagonal of $C_2 \times C_2$.

Lemma 9.10. *Applying the forgetful functor* $\mathrm{Mod}_{\mathsf{Cat}_\infty}(C)^{hC_2} \to \mathsf{Cat}_\infty^{hC_2}$ *after the functor* $\widetilde{T}_C$ *yields the functor* T_C*.*

Proof. This follows by construction. □

We denote by $\widetilde{sq}_C$ the composition of

$$m_{\mathrm{Mod}_{\mathsf{Cat}_\infty}(C)^{hC_2}}(2) \colon \mathrm{Mod}_{\mathsf{Cat}_\infty}(C)^{hC_2} \to \mathrm{Mod}_{\mathsf{Cat}_\infty}(C)^{hC_2}[C_2]$$

with $t_{\mathrm{Mod}_{\mathsf{Cat}_\infty}(C)}$.

Lemma 9.11. *We have a natural equivalence of functors* $\widetilde{T}_C \circ sq_C \simeq \widetilde{sq}_C \circ \widetilde{T}_C$*.*

Proof. This follows from the commutativity of the diagram (where we write $\mathrm{Mod}(C)$ for $\mathrm{Mod}_{\mathsf{Cat}_\infty}(C)$)

$$\begin{array}{ccccc}
\mathrm{Pic}(C)^{hC_2} & \longrightarrow & \mathrm{Mod}(C)^{h(C_2 \times C_2)} & \longrightarrow & \mathrm{Mod}(C)^{hC_2} \\
\downarrow & & \downarrow & & \downarrow \\
\mathrm{Pic}(C)^{h(C_2 \times C_2)} & \longrightarrow & \mathrm{Mod}(C)^{h(C_2 \times C_2 \times C_2)} & \longrightarrow & \mathrm{Mod}(C)^{h(C_2 \times C_2)} \\
\downarrow & & \downarrow & & \downarrow \\
\mathrm{Pic}(C)^{hC_2} & \longrightarrow & \mathrm{Mod}(C)^{h(C_2 \times C_2)} & \longrightarrow & \mathrm{Mod}(C)^{hC_2},
\end{array}$$

where the upper vertical maps are multiplication by 2 maps (where we keep track of the induced additional C_2-actions). □

Let D be a complete preadditive ∞-category and $X \in D[C_2]$. We denote by sq'_X the composition

$$X^{hC_2} \to (X^{hC_2})^{BC_2} \simeq X^{h(C_2 \times C_2)} \to X^{hC_2},$$

where the first map is $m_{X^{hC_2}}(2)$, $C_2 \times C_2$ acts on X via the first projection and the last map in the composition is induced via pullback along the diagonal of $C_2 \times C_2$.

Proposition 9.12. *Let the situation be as above. Then there is a natural commutative diagram*

$$\begin{array}{ccc}
X^{hC_2} & \xrightarrow{sq'_X} & X^{hC_2} \\
\downarrow & \nearrow_{\varphi} & \\
X & &
\end{array}$$

in D, where the vertical map is the natural map from the homotopy fixed points to X, and the composition of φ with this map $X^{hC_2} \to X$ is equivalent to the composition

$$X \xrightarrow{(\mathrm{id},\iota)} X \oplus X \to X,$$

where ι is the action of the nontrivial element of C_2 and the second map is the sum map.

Proof. The switching action of C_2 on $X \oplus X$ canonically exhibits $X \oplus X$ as object of $D[C_2][C_2] \simeq D[C_2 \times C_2]$. Restricting this action to the diagonal of $C_2 \times C_2$ gives rise to an object $\widetilde{X \oplus X} \in D[C_2]$. Letting C_2 act via the second factor of $C_2 \times C_2$ on $X \oplus X$ defines $(X \oplus X)' \in D[C_2]$. Note that there is a natural equivalence

$$(X \oplus X)' \simeq \widetilde{X \oplus X}$$

in $D[C_2]$ (using that $(X \oplus X)'$ is the free C_2-object on X). The claims of the Proposition then follow from the commutativity of the diagram

$$\begin{array}{ccccc}
X^{hC_2} & \longrightarrow & (X^{hC_2} \oplus X^{hC_2})^{hC_2} & \longrightarrow & (X^{hC_2})^{hC_2} \\
\downarrow & & \downarrow \simeq & & \downarrow \simeq \\
X & & (X \oplus X)^{h(C_2 \times C_2)} & \longrightarrow & X^{h(C_2 \times C_2)} \\
\downarrow \simeq & & \downarrow & & \downarrow \\
((X \oplus X)')^{hC_2} & \xrightarrow{\simeq} & (\widetilde{X \oplus X})^{hC_2} & \longrightarrow & X^{hC_2}
\end{array}$$

(the left square commutes since it commutes after mapping the right lower corner of this square to X via the projection to the first X-summand). □

Proof of Theorem 9.6. We specialize Proposition 9.12 to $D = \widetilde{\mathsf{SymMonCat}}_\infty$ and $X = \mathrm{Mod}_{\mathsf{Cat}_\infty}(C)$, where the C_2-action on X is the one described after Lemma 9.9. Then $sq'_X \simeq \widetilde{sq}_C$. Combining Lemmas 9.10 and 9.11 we see that

$$T_C \circ sq_C \simeq v \circ \widetilde{sq}_C \circ \widetilde{T}_C,$$

where v is the forgetful functor $\mathrm{Mod}_{\mathsf{Cat}_\infty}(C)^{hC_2} \to \mathsf{Cat}_\infty^{hC_2}$.

But for $L \in \mathrm{Pic}(C)^{hC_2}$ the image of $\widetilde{T}_C(L)$ under the functor $X^{hC_2} \to X$ is canonically equivalent to the tensor unit C, hence by Proposition 9.12 $\widetilde{sq}_C(\widetilde{T}_C(L))$ is also canonically equivalent to the tensor unit C, showing the claim. □

Remark 9.13. *Let the situation be as in Theorem 9.6. Then the underlying ∞-categories of the objects $T_C(sq_C(L))$ and $T_C(\mathbf{1})$ in $\mathsf{Cat}_\infty^{hC_2}$ are both canonically equivalent to C, thus the equivalence $T_C(\mathbf{1}) \simeq T_C(sq_C(L))$ in $\mathsf{Cat}_\infty^{hC_2}$ gives rise to an equivalence $\phi\colon C \to C$ in Cat_∞. We claim that ϕ is informally given by the assignment $C \ni X \mapsto X \otimes L$: Indeed, analyzing the proof of Proposition 9.12 we see that $\varphi\colon \mathrm{Mod}_{\mathsf{Cat}_\infty}(C) \to \mathrm{Mod}_{\mathsf{Cat}_\infty}(C)^{hC_2}$ sends the tensor unit C in $\mathrm{Mod}_{\mathsf{Cat}_\infty}(C)$ to $W := C \otimes_C \tau(C) \simeq \tau(C)$ together with a certain fixed point datum, where τ denotes the action of the nontrivial element of C_2 on $\mathrm{Mod}_{\mathsf{Cat}_\infty}(C)$. Also, after forgetting fixed point data, $\widetilde{sq}_C(\widetilde{T}_C(L))$ is $W' := C \otimes_C C$, and the identification between W' and W given by Proposition 9.12 is given by* $\mathrm{id}_C \otimes_C g$,

where g is informally given by $C \ni X \mapsto f(X)^\vee \otimes L$ (f the given autoequivalence of C). But after forgetting fixed point data the identification between the tensor unit of $\mathsf{Cat}_\infty^{hC_2}$ and W is given by $C \ni X \mapsto f(X)^\vee$, and composing the two identifications shows the claim.

Remark 9.14. *Let the situation be as in Remark 9.4. For $L \in \mathrm{Pic}(C)$ we have $sq_C(L) \simeq (L^{\otimes 2}, a) \in \mathrm{Pic}(C) \times A \simeq \mathrm{Pic}(C)[C_2]$ (where we view L in a canonical way as an object of $\mathrm{Pic}(C)[C_2]$) for an object $a \in A$. Then we see that $T_C(L^{\otimes 2}) \simeq T_C((\mathbf{1}, a^{\otimes(-1)}))$.*

Remark 9.15. *Let the situation be as in Corollary 9.3. Then for $L \in \mathrm{Pic}(C)^{hC_2}$ the spectra $T_C^{\mathrm{Sp}}(\mathbf{1})$ and $T_C^{\mathrm{Sp}}(sq_C(L))$ are canonically equivalent (even as $T_C^{\mathrm{Sp}}(\mathbf{1})$-module spectra).*

Remark 9.16. *Let the situation be as in Theorem 9.6. Proposition 9.12 shows that the functor $sq_C\colon \mathrm{Pic}(C)^{hC_2} \to \mathrm{Pic}(C)^{hC_2}$ also factors through $\mathrm{Pic}(C)$, in particular if the underlying object in $\mathrm{Pic}(C)$ of an object $L \in \mathrm{Pic}(C)^{hC_2}$ is trivial then so is $sq_C(L)$ (i.e. equivalent to the tensor unit).*

Remark 9.17. *Let the situation be as in Theorem 9.6. Then Remark 9.14 has the following generalization: For $L \in \mathrm{Pic}(C)^{hC_2}$ we have*

$$T_C(L^{\otimes 2}) \simeq T_C(L^{\otimes 2} \otimes sq_C(L)^{\otimes(-1)})$$

(note that $L^{\otimes 2} \otimes sq_C(L)^{\otimes(-1)}$ has trivial underlying object in $\mathrm{Pic}(C)$).

10 C_2-equivariant K-theory

When we do not take C_2-homotopy fixed points at the second stage in the definition of the functor K_h in section 5 we obtain a functor

$$K^{C_2}\colon \mathsf{SymMonCat}_\infty^{hC_2} \xrightarrow{(-)^\sim} \mathrm{Mon}_{E_\infty}(\mathsf{Spc})[C_2] \to \mathrm{Grp}_{E_\infty}(\mathsf{Spc})[C_2] \to \mathrm{Sp}[C_2]$$

which is similar to Proposition 5.1 lax symmetric monoidal. Forgetting the C_2-action in the target is the (direct sum) K-theory of the given symmetric monoidal ∞-category (forgetting its duality).

Let $\mathcal{O}$ be the opposite of the orbit category of C_2, so $\mathcal{O}^{\mathrm{op}}$ has two objects b and s, b has C_2 as monoid of self-maps, there is exactly one map from b to s and s has only the identity as self-map.

We can combine K^{C_2} and K_h to a lax symmetric monoidal functor

$$\begin{aligned} K_h^{C_2}\colon \mathsf{SymMonCat}_\infty^{hC_2} \xrightarrow{(-)^\sim} \mathrm{Mon}_{E_\infty}(\mathsf{Spc})[C_2] \to \mathrm{Mon}_{E_\infty}(\mathsf{Spc})^{\mathcal{O}} \\ \to \mathrm{Grp}_{E_\infty}(\mathsf{Spc})^{\mathcal{O}} \to \mathrm{Sp}^{\mathcal{O}}, \end{aligned}$$

where the second functor sends a BC_2-diagram in $\mathrm{Mon}_{E_\infty}(\mathsf{Spc})$ to the limiting cone (which is uniquely determined up to a contractible space of choices) of the diagram.

Remark 10.1. *The symmetric monoidal ∞-category $\mathrm{Sp}^{\mathcal{O}}$ is equivalent to the symmetric monoidal ∞-category of C_2-equivariant spectra indexed on the trivial universe.*

Corollary 10.2. *There is a natural functor*

$$\mathsf{SymMonCat}_\infty^{\mathrm{rig,preadd}} \to E_\infty(\mathrm{Sp}^{\mathcal{O}})$$

combining the multiplicative direct sum K-theory and hermitian K-theory functors.

For any $C \in \mathsf{SymMonCat}_\infty^{\mathrm{rig,preadd}}[C_2]$ the results of sections 8 and 9 show there is a lax symmetric monoidal functor

$$\mathrm{Pic}(C)^{hC_2} \to \mathrm{Sp}^{\mathcal{O}}.$$

11 Hermitian objects

Let Cat^1 be the 1-category of small categories and $\widehat{\mathsf{Cat}}^1$ the 1-category of not necessarily small categories. We view Cat^1 as object of $\widehat{\mathsf{Cat}}^1$.

Note there are natural functors $S\colon \mathrm{N}\mathsf{Cat}^1 \to \mathsf{Cat}$ and $\widehat{S}\colon \mathrm{N}\widehat{\mathsf{Cat}}^1 \to \widehat{\mathsf{Cat}}$.

Similarly to S we have the map $S'\colon \widehat{S}(\mathsf{Cat}^1) \to \mathsf{Cat}$ in $\widehat{\mathsf{Cat}}_\infty$.

Lemma 11.1. *Let G be a group. Let C be a (1-) category with G-action (in the strict sense, i.e. an object of $\widehat{\mathsf{Cat}}^1[G] = \mathrm{Fun}(BG, \widehat{\mathsf{Cat}}^1)$). Then there is a natural equivalence*

$$\widehat{S}(\mathrm{Fun}^G(EG, C)) \simeq \widehat{S}(C)^{hG}$$

in $\widehat{\mathsf{Cat}}$ (here EG is the translation groupoid of G).

We equip Cat^1 with the strict C_2-action given by $\mathsf{Cat}^1 \ni C \mapsto C^{\mathrm{op}}$. This induces a C_2-action on $\widehat{S}(\mathsf{Cat}^1)$.

Lemma 11.2. *The map S' is compatible with the C_2-actions.*

Proof. This follows from the fact that the nerve functor $\mathrm{N}\colon \mathsf{Cat}^1 \to \mathsf{sSet}$ is strictly compatible with the C_2-actions. □

Note $I \cong EC_2$ as C_2-categories. We get induced maps

$$\widehat{S}(\mathrm{Fun}^{C_2}(I, \mathsf{Cat}^1)) \simeq \widehat{S}(\mathsf{Cat}^1)^{hC_2} \to \mathsf{Cat}^{hC_2} \to \mathsf{Cat}_\infty^{hC_2}$$

in $\widehat{\mathsf{Cat}}_\infty$ (here we use that $\widehat{\mathsf{Cat}}$ is a full subcategory of $\widehat{\mathsf{Cat}}_\infty$), where the first equivalence is given by Lemma 11.1 and the second map is induced because of Lemma 11.2.

This composition can be equivalently viewed as a functor

$$\epsilon\colon \mathrm{N}\mathrm{Fun}^{C_2}(I, \mathsf{Cat}^1) \to \mathsf{Cat}_\infty^{hC_2}$$

of ∞-categories.

Remark 11.3. *It can be shown that the functor*

$$\mathrm{N}\mathrm{Fun}^{C_2}(I, \mathsf{Cat}^1) \to \mathsf{Cat}^{hC_2}$$

is a localization (in the naive sense, e.g. this functor has not a fully faithful right adjoint) at those natural transformations which are objectwise equivalences.

Lemma 11.4. $\mathrm{Fun}^{C_2}(I, \mathsf{Cat}^1)$ *is (isomorphic to) the category of small categories with strict duality and duality preserving functors in the sense of [15, section 2.1].*

Let $\triangle$ be the usual simplex category and

$$Q^{\mathrm{lax}}\colon \triangle \to \mathrm{Fun}^{C_2}(I, \mathsf{Cat}^1)$$

be the functor which sends $[n]$ to $[n] \star [n]^{\mathrm{op}}$, where $[n] \star [n]^{\mathrm{op}}$ is endowed with the unique strict duality.

Also for any $n \in \mathbb{N}$ we let Q_n be the category with strict duality obtained by forcing the maps $v_i\colon i \to i'$ in $[n] \star [n]^{\mathrm{op}}$ for any $0 \leq i \leq n$ to be identities (here i' is the object i viewed as object in $[n]^{\mathrm{op}}$).

There are projection maps $[n] \star [n]^{\mathrm{op}} \to Q_n$ in $\mathrm{Fun}^{C_2}(I, \mathsf{Cat}^1)$ which are localizations along the set of maps $\{v_i \mid 0 \leq i \leq n\}$ and which are part of a natural transformation between the functor Q^{lax} and a unique functor

$$Q\colon \triangle \to \mathrm{Fun}^{C_2}(I, \mathsf{Cat}^1)$$

with the property $Q([n]) = Q_n$ for any $n \in \mathbb{N}$.

We let

$$\widetilde{Q}^{\mathrm{lax}}\colon \mathrm{N}\triangle \to \mathsf{Cat}_\infty^{hC_2}$$

and

$$\widetilde{Q}\colon \mathrm{N}\triangle \to \mathsf{Cat}_\infty^{hC_2}$$

be the prolonguations of the functors $\mathrm{N}Q^{\mathrm{lax}}$ and $\mathrm{N}Q$ along ϵ.

The functors $\widetilde{Q}^{\mathrm{lax}}$ and $\widetilde{Q}$ induce canonical left adjoints

$$\widetilde{\mathcal{G}}^{\mathrm{lax}}\colon \mathsf{Spc}^{(\mathrm{N}\triangle)^{\mathrm{op}}} \to \mathsf{Cat}_\infty^{hC_2}$$

and

$$\widetilde{\mathcal{G}}\colon \mathsf{Spc}^{(\mathrm{N}\triangle)^{\mathrm{op}}} \to \mathsf{Cat}_\infty^{hC_2}.$$

We want to show that these functors factor through the localization

$$\mathsf{Spc}^{(\mathrm{N}\triangle)^{\mathrm{op}}} \to \mathsf{Cat}_\infty$$

(viewing Cat_∞ as complete Segal spaces).

Lemma 11.5. *The functors $\widetilde{Q}^{\mathrm{lax}}$ and $\widetilde{Q}$ preserve the standard colimits*

$$[1] \sqcup_{[0]} [1] \sqcup_{[0]} \dots \sqcup_{[0]} [1] \cong [n],$$

$n \in \mathbb{N}$, in $\triangle$, where on the left n times the object $[1]$ appears.

Thus the functors $\widetilde{\mathcal{G}}^{\mathrm{lax}}$ and $\widetilde{\mathcal{G}}$ factor through the localization

$$\mathsf{Spc}^{(\mathrm{N}\triangle)^{\mathrm{op}}} \to \mathsf{SegSpc}$$

of Segal spaces.

Lemma 11.6. *The functors $\widetilde{\mathcal{G}}^{\mathrm{lax}}$ and $\widetilde{\mathcal{G}}$ factor through complete Segal spaces.*

Proof. Since the functor $\mathsf{Cat}_\infty^{hC_2} \to \mathsf{Cat}_\infty$ detects equivalences it is sufficient to show that the induced compositions

$$G^{\mathrm{lax}}, G\colon \mathsf{SegSpc} \to \mathsf{Cat}_\infty^{hC_2} \to \mathsf{Cat}_\infty$$

send local equivalences (for the localization corresponding to complete Segal spaces) to equivalences which is equivalent to the fact that the right adjoints to G^{lax} and G factor through complete Segal spaces. This can be checked directly. □

We therefore get induced functors

$$\mathcal{G}^{\mathrm{lax}}, \mathcal{G}\colon \mathsf{Cat}_\infty \to \mathsf{Cat}_\infty^{hC_2}$$

whose right adjoints we denote by $\mathcal{H}^{\mathrm{lax}}$ and $\mathcal{H}$. For $C \in \mathsf{Cat}_\infty^{hC_2}$ we think of $\mathcal{H}^{\mathrm{lax}}(C)$ resp. $\mathcal{H}(C)$ as the ∞-categories of lax hermitian resp. hermitian objects of C.

Remark 11.7. *i) For $C \in \mathsf{Cat}_\infty^{hC_2}$ the ∞-category $\mathcal{H}(C)$ is naturally a full subcategory of $\mathcal{H}^{\mathrm{lax}}(C)$.*

ii) If C is a category with strict duality then there are natural equivalences $\mathcal{H}^{\mathrm{lax}}(\epsilon(C)) \simeq N(C_h)$ and $\mathcal{H}(\epsilon(C)) \simeq N(C_h^{\mathrm{str}})$ of ∞-categories (for the notation C_h see [15], and C_h^{str} denotes the full subcategory of C_h on those hermitian objects X such that the map $X \to X^\vee$ is an isomorphism).

$\mathcal{G}^{\mathrm{lax}}$ is not symmetric monoidal (for the cartesian structures), e.g. it does not preserve the tensor units (i.e. the final objects).

Proposition 11.8. *The functor $\mathcal{G}$ is symmetric monoidal for the cartesian structures.*

Proof. The only thing which has to be checked is that $\mathcal{H}$ preserves the terminal objects and that the map

$$\mathcal{G}(\Delta^1 \times \Delta^1) \to \mathcal{G}(\Delta^1) \times \mathcal{G}(\Delta^1)$$

is an equivalence, which can be done by hand. □

Proposition 11.9. *The diagram*

$$\begin{array}{ccccc} \mathsf{Cat}_\infty^{hC_2} & \xrightarrow{(-)^\sim} & \mathsf{Spc}^{hC_2} & \xrightarrow{\sim} & \mathsf{Spc}[C_2] \\ \downarrow{\scriptstyle \mathcal{H}} & & & & \downarrow{\scriptstyle (-)^{hC_2}} \\ \mathsf{Cat}_\infty & & \xrightarrow{\qquad (-)^\sim \qquad} & & \mathsf{Spc} \end{array}$$

commutes up to homotopy.

12 Applications

Assigning to a commutative ring R its symmetric monoidal category of finitely generated projective R-modules yields functors

$$\mathsf{CRing} \to \mathsf{SymMonCat}^{\mathrm{rig,preadd}} \to \mathsf{SymMonCat}_\infty^{\mathrm{rig,preadd}}$$

which we can prolong with either of the functors of Corollaries 7.9 and 10.2 to obtain (hermitian) K-theory functors.

This recovers the usual connective hermitian K-theory of R (see Remark 11.7 ii) and the corresponding references in the introduction).

Similarly we can assign to an E_∞-ring spectrum R its ∞-category of finitely generated projective modules $\mathcal{P}(R)$ (which is the full subcategory of the ∞-category of R-modules $\mathrm{Mod}(R)$ which are retracts of modules of the form R^n for some $n \in \mathbb{N}$) which inherits a rigid symmetric monoidal structure from $\mathrm{Mod}(R)$, thereby being an object of $\mathsf{SymMonCat}_\infty^{\mathrm{rig,preadd}}$.

Thus, again by prolonging with the functor of Corollary 10.2, we obtain a functor

$$E_\infty(\mathrm{Sp}) \to E_\infty(\mathrm{Sp}^{\mathcal{O}})$$

which should give a reasonable hermitian K-theory at least if R is connective. The underlying K-theory spectrum is connective algebraic K-theory (thereby carrying a C_2-action) if R is connective ([8, Theorem 5]).

Remark 12.1. *If we use instead of $\mathcal{P}(R)$ all perfect R-modules we obtain (hermitian) K-theory which should be considered to have a too big 0-th homotopy group since triangles will not be split into sums (as one would wish). A better construction of hermitian and real K-theory in this case can be found in [17].*

Remark 12.2. *Given any stable symmetric monoidal presentable ∞-category C the full subcategory of dualizable objects C^d will lie in $\mathsf{SymMonCat}_\infty^{\mathrm{rig,preadd}}$ thus giving rise to an E_∞-hermitian K-theory spectrum. Since this is direct sum hermitian K-theory its 0-th homotopy group will likely be considered to be too big (cf. Remark 12.1). Nevertheless we can apply the above machinery e.g. to the dualizable objects in stable motivic homotopy categories or in categories of triangulated motives. If the base scheme is a field of characteristic 0 these dualizable objects are just the compact objects. These remarks also apply in the setting of [17].*

Let $C \in \mathsf{SymMonCat}_\infty^{\mathrm{rig}}[C_2]$ and suppose C is stable. Then C^{hC_2} is also stable, and we have the object $K := \mathbf{1}[1] \in \mathrm{Pic}(C)^{hC_2}$ at our disposal. The object K gives rise to the object $M := sq_C(K) \otimes K^{\otimes(-2)} \in \mathrm{Pic}(C)^{hC_2}$. If C has trivial C_2-action then $\mathrm{Pic}(C)^{hC_2} \simeq \mathrm{Pic}(C) \times A$ and M can be considered as an object of A.

In particular for $R \in E_\infty(\mathrm{Sp})$ we can specialize to $C = \mathrm{Perf}(R)$ (perfect R-modules). Note that if we write $\mathrm{Pic}(\mathrm{Perf}(R))[C_2] \simeq \mathrm{Pic}(\mathrm{Perf}(R)) \times A$ then we have $\mathrm{Pic}(\mathcal{P}(R))[C_2] \simeq \mathrm{Pic}(\mathcal{P}(R)) \times A$, and we have the object $M \in A$ (M can be viewed as a C_2-enhancement of the automorphism "multiplication by -1" on the tensor unit of $\mathcal{P}(R)$).

Thus we can consider $T_{\mathcal{P}(R)}(M) \in (\mathsf{Cat}_\infty^{\mathrm{preadd}})^{hC_2}$ and $T_C^{\mathrm{Sp}}(M) \in \mathrm{Sp}$ ($T_C^{\mathrm{Sp}}(M)$ has also the structure of a $T_C^{\mathrm{Sp}}(\mathbf{1})$-module).

If we let R vary the objects M are compatible, thereby we get a functor

$$E_\infty(\mathrm{Sp}) \to \mathrm{Sp}$$

which can be considered as symplectic (hermitian) K-theory.

Remark 12.3. *In principle one can also consider tensor powers $M^{\otimes n}$, $n \in \mathbb{Z}$, of M. If $R \in \mathsf{CRing}$ then M has order at most 2 which recovers the known periodicity of hermitian K-theory. If R is n-truncated then it follows at least that the order of M is of the form 2^k for $0 \leq k \leq n+1$.*

Remark 12.4. *Let F be the free grouplike E_∞-space on the point (so F corresponds to the sphere spectrum) and view it as a Picard groupoid. If we write $F[C_2] \simeq F \times A$ then $\pi_0 A \cong \mathbb{Z}_2$, see [7, Theorem 1.1].*

Let $1 \in F$ be the object which corresponds to $1 \in \pi_0\mathbb{S}$, $\mathbb{S}$ the sphere spectrum. Then the A-component of $sq_F(1)$ is a non-zero element in $\mathbb{Z}_2$, in particular it has infinite order.

References

[1] Clark Barwick. On the algebraic k-theory of higher categories. *Journal of Topology*, 2016. doi:10.1112/jtopol/jtv042.

[2] Clark Barwick and Christopher Schommer-Pries. On the Unicity of the Homotopy Theory of Higher Categories. arXiv:1112.0040.

[3] Andrew J. Blumberg, David Gepner, and Gonçalo Tabuada. A universal characterization of higher algebraic K-theory. *Geom. Topol.*, 17(2):733–838, 2013.

[4] David Gepner, Moritz Groth, and Thomas Nikolaus. Universality of multiplicative infinite loop space machines. *Algebr. Geom. Topol.*, 15(6):3107–3153, 2015.

[5] Lars Hesselholt and Ib Madsen. Real algebraic k-theory. available at http://www.math.ku.dk/ larsh/papers/s05/.

[6] Max Karoubi. Théorie de Quillen et homologie du groupe orthogonal. *Ann. of Math. (2)*, 112(2):207–257, 1980.

[7] Wen Hsiung Lin. On conjectures of Mahowald, Segal and Sullivan. *Math. Proc. Cambridge Philos. Soc.*, 87(3):449–458, 1980.

[8] Jacob Lurie. Course notes for Math 281, Lecture 19. available at http://www.math.harvard.edu/ lurie/.

[9] Jacob Lurie. Higher Algebra. available at http://www.math.harvard.edu/ lurie/.

[10] Jacob Lurie. *Higher topos theory*, volume 170 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2009.

[11] Manos Lydakis. Smash products and Γ-spaces. *Math. Proc. Cambridge Philos. Soc.*, 126(2):311–328, 1999.

[12] Daniel Quillen. Higher algebraic K-theory. I. In *Algebraic K-theory, I: Higher K-theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972)*, pages 85–147. Lecture Notes in Math., Vol. 341. Springer, Berlin, 1973.

[13] Marco Schlichting. Hermitian k-theory, derived equivalences and Karoubi's fundamental theorem. available at http://homepages.warwick.ac.uk/ masiap/research/gwdg6Revised1.pdf.

[14] Marco Schlichting. Hermitian K-theory on a theorem of Giffen. *K-Theory*, 32(3):253–267, 2004.

[15] Marco Schlichting. The Mayer-Vietoris principle for Grothendieck-Witt groups of schemes. *Invent. Math.*, 179(2):349–433, 2010.

[16] Stefan Schwede. Stable homotopical algebra and Γ-spaces. *Math. Proc. Cambridge Philos. Soc.*, 126(2):329–356, 1999.

[17] Markus Spitzweck. A Grothendieck-Witt space for stable infinity categories with duality. Preprint.

[18] Bertrand Toën. Vers une axiomatisation de la théorie des catégories supérieures. *K-Theory*, 34(3):233–263, 2005.

[19] Friedhelm Waldhausen. Algebraic K-theory of topological spaces. I. In *Algebraic and geometric topology (Proc. Sympos. Pure Math., Stanford Univ., Stanford, Calif., 1976), Part 1*, Proc. Sympos. Pure Math., XXXII, pages 35–60. Amer. Math. Soc., Providence, R.I., 1978.

Fakultät für Mathematik, Universität Osnabrück, Germany.
e-mail:
haheine@Uni-Osnabrueck.DE
alopezavila@Uni-Osnabrueck.DE
markus.spitzweck@uni-osnabrueck.de